



**Section 24(c) Special Local Need Label**

**RESTRICTED USE PESTICIDE  
(GROUND AND SURFACE WATER CONCERNS)**

**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED  
APPLICATOR'S OR PERSONS UNDER THEIR DIRECT  
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE  
CERTIFIED APPLICATOR'S CERTIFICATION.**

**THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO  
GROUND AND SURFACE WATER CONCERNS. USERS MUST READ  
AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS  
FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE  
TO REACH GROUND AND SURFACE WATER.**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF  
OKLAHOMA**

**AATREX® 4L**

**EPA Reg. No. 100-497  
EPA SLN No. OK-830003**

**For Control Of Annual Weeds, Using AAtrex 4L In Tank Mixtures,  
During The Idle Season Following Winter Wheat Harvest In Early  
Summer Until Planting Back To Winter Wheat That Fall.**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL  
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

---

**DIRECTIONS FOR USE**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA registered AAtrex 4L label and on the labels of herbicides tank mixed with AAtrex.

This labeling must be in the possession of the user at the time of pesticide application.

**DO NOT** contaminate or apply to ponds, lakes, or streams.

AAtrex is a broad spectrum herbicide which, in tank mixture with an EPA-registered herbicide for use in fallow, is effective for preemergence control of carpetweed, common lambsquarters, common purslane, kochia, mustards, Russian thistle, tumble and other pigweeds, wild lettuce, and suppression of volunteer wheat.

For best results, apply herbicide after straw has been uniformly distributed over the wheat stubble or soil surface.

Winter Wheat - After Harvest for Weed Control During Idle Season Until Fall Planting to Winter Wheat

**Precautions:** **DO NOT** apply on soils having a pH greater than 7. **DO NOT** apply in the Oklahoma Panhandle. **DO NOT** treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil. **DO NOT** treat soils with calcareous surface layers. **DO NOT** use on sands. Avoid spray overlap.

Apply AAtrex 4L at 0.5-1 pt./A in combination with an EPA-registered herbicide for use in a 4-month fallow, as soon as possible following wheat harvest.

Use the lower rates on coarse-textured soils and soils relatively low in organic matter; use the higher rates on fine-textured soils and soils relatively high in organic matter.

For control of established weeds, tank mix with paraquat at 1-2 pts./A + X-77® at 1-2 pts. per 100 gals. of spray solution, or stubble plow to control large established weeds. Refer to the herbicide labels for directions, weeds controlled, precautions, and limitations.

Apply in a minimum of 20 gals. of water per acre with ground equipment or in a minimum of 5 gals. of water per acre by air.

If weeds or volunteer wheat become a problem, a shallow cultivation or a postemergence application of paraquat + X-77 should be employed.

**Note:** **DO NOT** feed green forage or graze treated areas to livestock for six months following application. Treat only once during the same fallow period. **DO NOT** plant wheat until four months after application.

**Chemigation:** **DO NOT** apply these products through any type of irrigation system.

AAtrex® trademark of Syngenta  
X-77® trademark of Loveland Industries, Inc.

24(c) registrant:  
Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, NC 27419-8300

EPA SLN No.: OK-830003

Label Code: OK0497024BA1103

Word: OKAAtrex4LWheatFallowSLNLbI9-17-03