

# SUPPLEMENTAL LABELING

This supplemental label expires on March 22, 2025 and must not be used after this date

## RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO NON-TARGET INVERTEBRATES, MAMMALS, AND AQUATIC ORGANISMS  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT  
SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.



ABAMECTIN GROUP 6 INSECTICIDE

EPA Reg. No. 89118-10

### COTTON IN-FURROW APPLICATIONS

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of the application. Read the label affixed to the container for Averland FC before applying. Use of Averland FC according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Averland FC.

#### COTTON

SOIL PESTS	USE RATES	
	fl oz product/A (lb a.i./A)	fl oz product/1000 row ft (oz a.i./1000 row ft) @ 36" rows
Suppression: Nematodes	3.5 (0.019)	0.24 (0.021)

#### At-Plant / Soil Applied Instructions

For early season seed and seedling protection apply in the following method:

- Apply in-furrow in a minimum of 5 gallons per acre of final volume as a spray or dribble directed on the seed during planting.
- Linear application rate is based on 36" rows. For different row spacing consult the Linear Application Rate Conversion Chart in the Instructions for At-Plant

Applications section. In all cases do not exceed the maximum per acre use restriction.

#### Specific Use Restrictions:

- Not for use in California
- **Application Method:** Do not apply through any type of irrigation systems.
- **Maximum Single Application:** Do not apply more than 3.5 fl oz/A (0.019 lb a.i./A) per application.
- **Annual Maximum:**
  - Do not exceed 7.0 fl oz Averland FC/A/calendar year.
  - Do not exceed 0.038 lb abamectin/A/calendar year from all abamectin containing products including all application types (seed treatments, soil, foliar).
- **Application Interval:** Do not make applications less than 21 days apart.
- **Pre-Harvest Interval (PHI):** Do not apply within 20 days of harvest.
- **Resistance Management:** Do not make more than 2 foliar applications of all abamectin containing products per year.
- **Grazing:** Do not graze livestock in treated areas or feed treated foliage to livestock.



Vive Crop Protection, Inc.  
500 Westover Dr #10198  
Sanford, NC 27330  
1-888-760-0187

Averland, Allosperse, and Vive Crop Protection are trademarks of Vive Crop Protection Inc.

© 2022 Vive Crop Protection Inc. All rights reserved.