



EPA REG. N.º: 91473-1-86182
RECOMMENDATION FOR THE CONTROL OF
ERWINIA CAROTOVORA/BACTERIAL SOFT ROT/AERIAL STEM ROT

FOR DISTRIBUTION AND USE IN ALL STATES EXCEPT California

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted by Section 2 (ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Recommendation for the control of Erwinia carotovora/bacterial soft/aerial stem rot: This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the applicable directions, restrictions and precautionary statements plus all the other information appearing on the EPA-registered product label.

Crop	Pest	Rate (oz/acre)	Timing/Application
Root/Tuber and Corm Vegetables Carrot Potato Sweet Potato Beets Ginger Horseradish Radish Ginseng Turnip and other root/tuber and corm crops including those grown for seed production	Black leg/bacterial soft rot/Aerial Stem Rot (Erwinia carotovora)	20-30	For suppression, begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat on a 7- to 10-day interval or as needed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. For use only as described on this label.

MANUFACTURED BY:

Seipasa, S.A.
22240 Tardienta (Huesca) Spain
www.seipasa.com

DISTRIBUTED BY:

Stockton (Israel) Ltd
17, Ha Mefalsim str., Petach Tikva
4951447, Israel
Tel In USA: +1-833-551-CROP (2767)
www.stk-ag.com