

---

**PRODUCT INFORMATION**

---



**Syngenta Crop Protection, Inc.**  
Greensboro, NC 27409  
www.syngenta-us.com

**SUPPLEMENTAL LABEL FOR ABOUND®**

**FUNGICIDE**

**EPA Reg. No. 100-1098**

Grasses grown for seed removal of feeding restriction

Active Ingredient:

Azoxystrobin: methyl ( <i>E</i> )-2-{2-[6-(2-cyanophenoxy) pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate* .....	22.9%
Other Ingredients:	77.1%
Total:	100.0%

Contains 2.08 lbs. of active ingredient per gallon

\*IUPAC

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**Before using Abound as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

## DIRECTIONS FOR APPLICATIONS

Crop	Target Diseases	Use Rate fl oz./product/A	Remarks
<b>Grasses</b> (grown for seed)	Rust ( <i>Puccinia</i> spp.)  Powdery mildew  ( <i>Erysiphe graminis</i> ) Ergot Stem Diseases	6.2-15.4 (0.10-0.25)	<p><b><u>Integrated Pest (Disease) Management:</u></b> Abound should be integrated into an overall disease management strategy that includes varieties with disease tolerance, proper timing of irrigation, crop rotation, and fertility.</p> <p><b><u>Resistance Management:</u></b> Do not apply more than two sequential foliar applications of Abound or other strobilurins before alternating with a fungicide that has a different mode of action. Do not make more than two (2) applications of Abound per acre per year.</p> <p><b><u>Application Directions:</u></b> Abound applications should begin prior to disease development and continue throughout the season on a 10-14 day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at recommended rates.</p>
<p><b>Specific Use Restrictions:</b> Do not apply more than 1.44 quarts (0.75 lb. active ingredient.) per acre per season.                      May be applied up to 8 days prior to harvest (swathing).</p>			

Trademark of Syngenta

©2004 Syngenta

Syngenta Crop Protection, Inc.  
 Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP 1098A-S9 0304

ABO 1098A-S9 0304 – nsp 03-30-04