

ACCELERON[®]

I-609

Insecticide Seed Treatment

Seed treatment use on corn and sorghum for the control of listed insect pests.
All seed treated with this product must be conspicuously colored at the time of treatment.

Active Ingredient:

clothianidin	48.0%
--------------------	--------------

Other Ingredients:	52.0%
---------------------------------	--------------

Total:	100.0%
---------------------	---------------

Contains 5.00 lbs clothianidin per U S gallon

EPA Reg. No. 7969-458-264

EPA Est. No. is indicated by the letter(s) immediately following the first five numbers of the lot code printed on the container. 5905-GA-001 (L); 51036-GA-001 (S)

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUTION

Please refer to booklet for additional precautionary statements and directions for use.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours a Day 1-800-334-7577.

For Product Information Call 1-866-99BAYER (1-866-992-2937)

Net Contents: 2.5 GAL

191101C 08/21
13186V1-11

PRODUCED FOR: Bayer CropScience LP
800 N. Lindbergh Blvd. • St. Louis, Missouri 63167

PRODUCT OF GERMANY

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told to by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance; then give artificial respiration, preferably mouth to mouth if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
In case of emergency, call toll free the BAYER CROPSCIENCE Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor or going for treatment.	

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Harmful if absorbed through the skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeve shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Environmental Hazards

This product is toxic to birds and aquatic invertebrates. **DO NOT** apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. **DO NOT** apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on the label.

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spill, or equipment leak and dispose of wastes according to the disposal instructions on this label.

Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect spilled seeds. This compound is highly toxic to bees exposed directly (contact).

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeve shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber \geq 14 mils, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride rubber \geq 14 mils or viton \geq 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

DO NOT contaminate water, food, or feed by storage or disposal. Store in a cool, dry secured storage area.

Pesticide Disposal

To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Plastic Containers

Rigid nonrefillable containers (equal to or less than 5 gallons)

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration.

Product Information

Acceleron I-609 Insecticide Seed Treatment

- Systemic insecticide seed treatment for use on corn and sorghum for the control of listed insects.
- Provides early season protection against the listed insect pests.
- Provides protection under low-to-moderate corn rootworm pressure.

Application Instructions

Acceleron I-609 Insecticide Seed Treatment

- To be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time. Pretest all tank mixes to determine physical compatibility between formulations. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.
- Treatment of highly mechanically damaged seed, or seed of known low vigor or poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of BAYER CROPSCIENCE, no claims are made to guarantee germination of carryover seed.

Use Restrictions

- All seed treated with this product must be conspicuously colored at the time of treatment. This product contains no colorant. An EPA-approved dye or colorant must be added when this product is applied to seed to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals. Regulations pertaining to coloration of treated seed enforced by 40 CFR 153.155 must be strictly adhered to when using this product.
- For use only in commercial seed treatment facilities. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.
- Regardless of method of application (seed treatment, soil, or foliar), **DO NOT** apply more than 0.2 lb of active ingredient clothianidin per acre per year.
- Not for use on sorghum seed in the state of California.

Refer to the specific use directions and restrictions in each crop table.

Specific Crop Directions

Commercial Seed Treatment Facilities

Corn (field corn, popcorn)	
Insect Pests	Product Rate fl oz/80,000 seeds (mg ai/seed)
Corn flea beetle (through first true seed leaf) Seedcorn maggot ² Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) White grubs ² (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae)	0.57 (0.125)
Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) Sugarcane beetle Thrips White grubs (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae) Wireworms	1.13 to 2.26 (0.25 to 0.50)
Corn rootworms (including Mexican, northern, southern, western) Southern corn billbug	5.64 (1.25)
Crop-specific Directions ¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed. ² For seed protection only.	

Corn (Sweet Corn)	
Insect Pests	Product Rate fl oz/80,000 seeds (mg ai/seed)
Corn flea beetle (through first true seed leaf) Seedcorn maggot ² Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) White grubs ² (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae)	0.57 (0.125)
Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) Sugarcane beetle Thrips White grubs (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae) Wireworms	1.13 to 2.26 (0.25 to 0.50)
Crop-specific Directions ¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed. ² For seed protection only.	

Sorghum	
Insect Pests	Product Rate fl oz/100 lbs seeds (mg ai/seed) ¹
Aphids (including greenbug, corn leaf aphid, English grain aphid, yellow sugarcane aphid) Black granulate cutworm Chinch bugs White grubs Wireworms	5.1 to 6.4 (0.05 to 0.06)
Crop-specific Restrictions • Not registered for use in California. ¹ Rate is calculated based on number of seeds/lb averages of sorghum (18,000).	

Rotational Crop Plant-Back Intervals					
Crop Planted	Immediate Plant-back	30-day Plant-back	4-month Plant-back	8-month Plant-back	12-month Plant-back
Field Corn, Popcorn	canola, cereal grains, cotton, soybean,	dry beans, grasses, non-grass animal feeds		Sugarcane	Any crop without an earlier plant-back interval
Sweet Corn, Sorghum	Brassica (Cole) vegetables, bulb vegetables, cucurbit vegetables, fruiting vegetables, leafy green vegetables, root and tuber vegetables		Any crop without an earlier plant-back interval		

SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with **Acceleron I-609 Insecticide Seed Treatment** must be labeled with the following statements:

- This seed has been treated with **Acceleron I-609 Insecticide Seed Treatment**, which contains **48.0% clothianidin**.

• **DO NOT** use treated seed for food, feed, or oil production.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with **Acceleron I-609 Insecticide Seed Treatment**:

- **Store treated seeds away from food and feedstuffs.**

• **DO NOT allow children, pets, or livestock to have access to treated seed.**

• **Wear long-sleeve shirt, long pants, and chemical-resistant gloves when handling treated seed.**

- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

• Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

• Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.

• **DO NOT** contaminate water bodies when disposing of planting equipment washwater.

• Dispose of seed packaging or containers in accordance with local requirements. **DO NOT** use empty seed bags for any other purpose.

• This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds.

• This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.

• **DO NOT** apply a neonicotinoid insecticide foliar spray after planting treated seed. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

• **Field corn, popcorn**

- o Immediate plant-back: canola, rapeseed, cereal grains, field corn, popcorn, sweet corn, root and tuber vegetables, soybean
- o 30 day plant-back: dry beans, grasses, non-grass animal feeds

- o 8 month plant-back: sugarcane

- o 12 month plant-back: any crops without an earlier plant-back interval

• **Sweet corn, sorghum**

- o Immediate plant-back: canola, rapeseed, cereal grains, field corn, popcorn, sweet corn, root and tuber vegetables, sorghum, soybean

- o 30 day plant-back: dry beans, grasses, non-grass animals, feeds

- o 4 month plant-back: any crop without an earlier plant-back interval

• Follow-up foliar sprays of registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.

• Regardless of method of application (seed treatment, soil, or foliar) **DO NOT** apply more than 0.2 lb of active ingredient clothianidin per acre per year.

• Not for use on sorghum seed in the state of California.

• **DO NOT** use **Acceleron I-609 Insecticide Seed Treatment** in areas where rice/crawfish aquaculture practices are in place, or in or near fish farms, shrimp, prawn or crab pond or nursery operations.

• **Corn (field corn, popcorn)** - Treated at the rate of XX mg ai/seed.

• **Corn for pest insect southern corn billbug** - Follow-up foliar sprays of a locally registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.

• **Corn (sweet)** - Treated at the rate of XX mg ai/seed.

• **Sorghum** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience (Bayer), no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER'S ELECTION, THE REPLACEMENT OF PRODUCT.

Acceleron® is a registered trademark of Bayer Group.

©2021 Bayer Group. All rights reserved.

NVA 2019-04-602-0106_8058CR7

ACCELERON[®]

I-609

Insecticide Seed Treatment

Seed treatment use on corn and sorghum for the control of listed insect pests. All seed treated with this product must be conspicuously colored at the time of treatment.

Active Ingredient:

clothianidin **48.0%**

Other Ingredients: **52.0%**

Total: **100.0%**

Contains 5.00 lbs clothianidin per U S gallon

EPA Reg. No. 7969-458-264

EPA Est. No. is indicated by the letter(s) immediately following the first five numbers of the lot code printed on the container. 5905-GA-001 (L); 51036-GA-001 (S)

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUTION

Please refer to booklet for additional precautionary statements and directions for use. **FIRST AID: If swallowed:** Immediately call a poison control center or doctor for treatment advice. **DO NOT** induce vomiting unless told to by a poison control center or doctor. Have person sip a glass of water if able to swallow. **DO NOT** give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or ambulance; then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **In case of emergency, call toll free the BAYER CROPSCIENCE Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor or going for treatment. NOTE TO PHYSICIAN:** No specific antidote is available. Treat the patient symptomatically. **Precautionary Statements: Hazards to Humans and Domestic Animals: CAUTION.** Harmful if swallowed. Harmful if absorbed through the skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. **Personal Protective Equipment (PPE)** Applicators and other handlers must wear: Long-sleeve shirt, Long pants, Chemical-resistant gloves made out of: barrier laminate, nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride rubber ≥ 14 mils, or viton ≥ 14 mils, and Shoes plus socks. **Environmental Hazards:** This product is toxic to birds and aquatic invertebrates. **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. **DO NOT** apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on the label. This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spill or equipment leak and dispose of wastes according to the disposal instructions on this label. Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect spilled seeds. This compound is highly toxic to bees exposed directly (contact). **STORAGE AND DISPOSAL: DO NOT** contaminate water, food or feed by storage, disposal, or cleaning of equipment. **Pesticide Storage: DO NOT** contaminate water, food, or feed by storage or disposal. Store in a cool, dry secured storage area. **Pesticide Disposal:** To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry). **Container Handling: Nonrefillable Plastic Containers: Rigid nonrefillable containers (equal to or less than 5 gallons):** Nonrefillable Container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Once container is rinsed, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration.

For MEDICAL AND TRANSPORTATION Emergencies ONLY Call 24 Hours a Day 1-800-334-7577. For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Acceleron[®] is a registered trademark of Bayer Group. ©2021 Bayer Group. All rights reserved.

NVA 2019-04-602-0106 8058CR7

191101C 08/21

Net Contents: 2.5 GAL

Product of Germany

13186V1-11

PRODUCED FOR: Bayer CropScience LP
800 N. Lindbergh Blvd. • St. Louis, Missouri 63167