

Section 2 (ee) Recommendation

Date: August 20, 2021

Product: Actigard® 50WG Plant Activator

EPA Reg. No. 100-922

Use: Enhanced Control of Neopestalotiopsis Leaf Spot and Fruit Rot in Strawberry

State: FL, GA, MD, NC, NJ, PA, SC, VA, WV

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

For enhanced control of Neopestalotiopsis Leaf Spot and Fruit Rot, apply Actigard 50WG at 0.375-0.75 oz/A in combination with an approved efficacious fungicide when disease first appears and continue on a 7 to 14 day interval. Do not apply within 5 days of transplanting.

Refer to the Crop Use Directions for Low Growing Berry, Crop Subgroup 13-07G, on the Actigard 50WG Plant Activator Section 3 label for other use directions, use restrictions and precautions.

Actigard® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2021 Syngenta

2(ee) Registrant: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419-8300

Label Code: FL0922104BC1023