



FIFRA 2ee Supplemental Labeling

ADVANTIGRO®

EPA Reg. No. 62097-43-2935

ACTIVE INGREDIENT:

Cytokinin (as kinetin)	0.009%
Indole Butyric Acid	0.005%
Gibberellic Acid (A ₃)	0.005%
OTHER INGREDIENTS	99.981%
Total	100%

KEEP OUT OF REACH OF CHILDREN

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the ADVANTIGRO® container before applying. Before using ADVANTIGRO in accordance to this Supplemental Label, follow all precautionary statements, restrictions, Worker Protection Standard requirements and applicable use directions on the label affixed to the product container.

DIRECTIONS FOR USE

Table 1. Crop Application Rates and Application Instructions: FIELD CROPS

Crop	No. of Applications	Rate (fl.oz./Acre) by Type of Application			Application Timing for Banded and Broadcast/Foliar
		In-Furrow	Banded	Broadcast Foliar	
Corn	1	2	2	2	Make application at 2-6 leaf stage.
	1		2	2	Apply during flowering from initiation of flowering to end of bloom stage.
Soybeans	1	2	2	2	Make application at V4-V8 (3-7 trifoliolate).
	2		2	2	Make application at V4-V8 (3-7 trifoliolate). Repeat applications can be made on 10-17 day intervals.

Table 2. Crop Application Rates and Application Instructions: VEGETABLE CROPS

Crop	No. of Applications	Rate (fl.oz./Acre) by Type of Application			Application Timing for Banded and Broadcast/Foliar
		In-Furrow	Banded	Broadcast Foliar	
Corn, sweet	1	2	2	2	Make application at 2-6 leaf stage.
	2 or more		2	2	Make first application at 2-6 leaf stage. Repeat applications can be made on 7-21 day intervals through end of tasseling.
Cucumbers	1	2	2	2	Make application between flower bud initiation and first bloom.
	2-3		2	2	Make first application between flower bud initiation and first bloom. Repeat applications can be made on 7-10 day intervals.
	3-4		2	2	Direct Seeded: First application at 3-4 leaf stage. Repeat application can be made on 7-10 day intervals.
Melons	2-3	2	2	2	Make first application at flower bud initiation. Repeat applications can be made on 7-10 day intervals.
	4-6		2	2	Make first application 2 weeks after emergence. Repeat applications can be made on 7-14 day intervals.
Potatoes	1	2	2	2	Apply at tuber initiation.
	3		2	2	Make first application at stolon formation (8-10 leaf stage). Repeat applications can be made on 10-14 day intervals.

WILBUR-ELLIS COMPANY LLC
 PO BOX 16458
 FRESNO, CA 93755
 (559) 442-1220

This Supplemental Label for ADVANTIGRO expires on December 31, 2019

WILBUR-ELLIS® logo and Ideas to Grow With® and ADVANTIGRO® are registered trademarks of Wilbur-Ellis Company LLC.