

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1324B-S3 1016**

**GROUP 3 11 FUNGICIDES**

**Aframe™ Plus**

Broad spectrum fungicide for control of plant diseases

**This supplemental label expires on October 21, 2019 and must not be used or distributed after this date.**

Active Ingredients:	
Azoxystrobin.....	13.5%
Propiconazole .....	11.7%
Other Ingredients:	74.8%
Total:	100.0%

Contains 1.02 lb ai propiconazole and 1.18 lb ai azoxystrobin per gallon

Aframe Plus is formulated as a suspo-emulsion (SE).

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**EPA Reg. 100-1324**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Aframe Plus as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

## DIRECTIONS FOR USE

<b>Rotational Crops</b>	<b>Planting Time From Last Aframe Plus Application</b>
Brassica, leafy greens (subgroup 5B) Bulb crops Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale) Corn (field, seed, popcorn, and sweet) Dill Grasses grown for seed (Cool season grasses only) Mint Oats Peanuts Quinoa Radish Rice Rye Sorghum Soybeans Strawberries Sugar beets Wild rice Watercress	0 days
Buckwheat Millet	12 Months
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/season)	75 days
All Other Crops Intended for Food and Feed	105 days

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Grasses grown For Seed</b> (Cool season grasses only)	Ergot Stem Diseases Powdery Mildew <i>(Erysiphe graminis)</i> Rusts <i>(Puccinia spp.)</i>	14 - 26	<p>Apply Aframe Plus when powdery mildew infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 26 fl oz/A (except bluegrass apply 14 fl oz/A) and make applications at 14-day intervals until the seed is mature. For bluegrass, it is important to begin application early in the growing season.</p> <p>Make no more than 2 sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.</p>

**Application:** Aframe Plus is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Apply Aframe Plus in a minimum of 20 gal of water per acre for ground or in a minimum of 10 gal of water per acre for aerial. Aframe Plus may be applied by ground, air, or chemigation.

**Specific Use Restrictions:**

- 1) Do not feed hay cut within 20 days of the last application.
- 2) Do not graze treated areas within 140 days of the last application.
- 3) Do not apply more than 86.0 fl oz/A/season of Aframe Plus.
- 4) Do not apply more than 0.90 lb ai propiconazole-containing products/A/season.
- 5) Do not apply more than 0.8 lb ai azoxystrobin-containing products/A/season.
- 6) Do not apply within 20 days of harvest (20-day PHI) of seed.
- 7) Do not apply to bermudagrass grown for seed.

<b>Crop</b>	<b>Target Diseases</b>	<b>Use Rate fl oz product/A</b>	<b>Remarks</b>
<b>Mint</b>  Peppermint Spearmint	Powdery Mildew ( <i>Erysiphe</i> spp.)  Rust ( <i>Puccinia menthae</i> )	10.5 - 14	Begin applications when the plants are 2-4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

**Application:** Aframe Plus may be applied by aerial, ground (minimum of 20 gal/A) or chemigation.

**Specific Use Restrictions:**

- 1) Do not apply more than 28 fl oz/A of Aframe Plus per crop.
- 2) Do not apply more than 0.22 lb ai of propiconazole-containing products/A/season.
- 3) Do not apply more than 0.75 lb ai of azoxystrobin-containing products/A/season.
- 4) Do not apply within 7 days of harvest (7-day PHI).

Aframe™ Plus and the Syngenta logo are trademarks of a Syngenta Group Company

©2017 Syngenta

Aframe Plus SCP 1324B-S3 1016 – ep – 4-28-17