



**FIFRA 24 (C)**  
**Special Local Need Label**  
**For Use on Hops Only**

EPA Reg. No. 279-3241      EPA SLN. No. MI-140010

## FIFRA Section 24 (c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MICHIGAN.**

**This label is valid until September 17, 2019 or until otherwise amended, withdrawn, canceled, or suspended.**

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

| CROP/SITE: | TARGET PEST:  | RATE:   | METHOD OF APPLICATION:  | FREQUENCY AND TIMING OF APPLICATION:   |
|------------|---|---|---|--|
| Hops       | Apply Aim EC using shielded sprayers or hooded sprayers to control emerged and actively growing broadleaf weeds within or between the rows of the crop. Refer to weed control list below. | 0.5 – 2.0 fl. oz. (0.008 – 0.032 pound active ingredient) per acre. Refer to weed control list below. | Apply Aim EC at 2.0 fl. oz. (0.03 pound active ingredient) per acre per application in a minimum of 20 gallons of spray solution by boom-type ground application equipment only to the basal portion of the hop plant (approximately the lower 1.5 feet) and to the sucker mat which extends from the base of the plant to approximately 1.5 to 2 feet into the row. An alternate row treatment program may be followed to avoid the removal of excessive photosynthetic capacity from the crown area. When treating alternating rows on different days, the equivalent maximum rate must not exceed 3.2 fl. oz. (0.015 pound active ingredient) of Aim EC per application per treated row area totaling 0.5 acres. | Do not apply more than 7.6 fl. oz. (0.12 pound active ingredient) per acre per season. Allow 14 days between treatments of Aim EC. |

**Use Restrictions:**

- Do not apply Aim EC using air blast or air assisted sprayers.
- Do not apply within 7 days of harvest.
- Do not apply through any type of irrigation system.

**This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.**

Aim is a trademark of FMC Corporation.  
 ©2014 All rights reserved.  
 09-19-14

**FMC**  
 FMC Corporation  
 Agricultural Solutions  
 1735 Market Street  
 Philadelphia, PA 19103

|  |   |
|--|---|
| <b>Weeds Controlled</b>  | <b>Aim EC Use Rate</b><br><b>fl. oz. (pound active ingredient) per acre</b> |
| Lambsquarters, common (up to 3 inches tall)  | 0.5 fl. oz. (0.008 pound active ingredient) per acre                        |
| Morningglory, ivyleaf (up to 3 leaves)   |   |
| Morningglory, pitted (up to 3 leaves)  |   |
| Nightshade, Eastern black  |   |
| Pigweed, redroot   |   |
| Velvetleaf   |   |
| Waterhemp (up to 2 inches tall)  |   |
| <b>Weeds Controlled</b>  | <b>Aim EC Use Rate</b><br><b>fl. oz. (pound active ingredient) per acre</b> |
| All the weeds controlled at 0.5 fl. oz. (0.008 pound active) per acre plus the weeds listed below: | 0.8 fl. oz. (0.013 pound active ingredient) per acre                        |
| Cheeseweed   |   |
| Filaree, redstem   |   |
| Flixweed   |   |
| Lambsquarters, common  |   |
| Morningglory, entireleaf   |   |
| Morningglory, ivyleaf  |   |
| Morningglory, pitted   |   |
| Morningglory, scarlet  |   |
| Nightshade, hairy  |   |
| Pennycress, field  |   |
| Pigweed, prostrate   |   |
| Pigweed, smooth  |   |
| Pigweed, tumble  |   |
| Purslane, common   |   |
| Sesbania, hemp   |   |
| Smartweed, PA (seedling)   |   |
| Spurge, prostrate  |   |
| Tansymustard   |   |
| Velvetleaf (24")   |   |
| Waterhemp, common & tall   |   |
| <b>Weeds Controlled</b>  | <b>Aim EC Use Rate</b><br><b>fl. oz. (pound active ingredient) per acre</b> |
| All the weeds controlled at 0.8 fl. oz. (0.013 pound active) per acre plus the weeds listed below: | 1.0 fl. oz. (0.016 pound active ingredient) per acre                        |
| Amaranth, spiny  |   |
| Anoda, spurred   |   |
| Bedstraw, catchweed  |   |
| Buffalobur   |   |
| Carpetweed   |   |
| Cocklebur  |   |
| Copperleaf, hophornbeam  |   |
| Cotton, GMO Varieties  |   |
| Cotton, volunteer  |   |
| Eclipta  |   |
| Fiddleneck, coast  |   |
| Groundcherry, smooth (seedling)  |   |
| Groundcherry, Wright's   |   |
| Jimsonweed   |   |
| Kochia   |   |
| Lettuce, Prickly 2-3 leaf  |   |
| Nettle, burning  |   |
| Nightshade, American black   |   |
| Nightshade, black  |   |
| Rocket, London   |   |
| Shepherdspurse   |   |
| Speedwell, Virginia  |   |
| Spiderwort, tropical   |   |
| Thistle, Russian   |   |
| Wallflower, bushy  |   |

| <b>Weeds Controlled</b>  | <b>Aim EC Use Rate<br/>fl. oz. (pound active ingredient) per acre</b> |
|--|---|
| All the weeds controlled at 1.1 fl. oz. (0.016 pound active) per acre plus the weeds listed below: | 1.6 fl. oz. (0.025 pound active ingredient) per acre                  |
| Amaranth, Palmer   |   |
| Corn Spurry  |   |
| Filaree, broadleaf   |   |
| Filaree, white   |   |
| Lettuce, prickly   |   |
| Mallow, Venice (up to 2 inches tall)   |   |
| Meadowfoam   |   |
| Redmaids   |   |
| <b>Weed List, Burndown of top growth</b>   |   |
| Bindweed, field  | 1.0 – 2.0 fl. oz. (0.016 – 0.032 pound active ingredient) per acre    |
| Burclover  |   |
| Dayflower  |   |
| Sage, lanceleaf  |   |
| Sowthistle   |   |

**\*When used as directed, Aim EC will provide control of the listed weeds up to four (4) inches in height, or as specified.**