

Alexin[®] 0-0-8

A potassium, calcium and magnesium true solution with PAC* chelation technology

GUARANTEED ANALYSIS:

Soluble Potash (K ₂ O)	8.0%
Calcium (Ca)	2.4%
2.4% Chelated Calcium	
Magnesium (Mg)	0.8%
0.8% Chelated Magnesium	

Derived from potassium hydroxide, calcium hydroxide and magnesium oxide chelated with PolyAldoCarbosate.

For use as a fertilizer in foliar and fertigation applications to supplement potassium, calcium and magnesium levels and to boost plant metabolism in agricultural crops.

***PAC refers to NutriAg's "PolyAldoCarbosate" Technology**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

DIRECTIONS FOR USE (Use only as Directed):

- Exercise caution with foliar applications in extreme weather conditions such as high temperatures and poor drying conditions.
- Aerial application – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 10 gallons of water/acre. Apply with sufficient water to deliver spray uniformly on leaf surfaces. Higher volumes of water should be used for higher rates of fertilizer.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- Alexin is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions.
- **Consult a NutriAg representative for specific crop recommendations, application timings and necessity of its use.**

SUGGESTED RATES

<p>FOLIAR RATES: For crop remediation, apply Alexin before and after each stress event at 1 qt/ac, 7-14 days apart. Alternatively for general plant health, use 1 pt/acre at approximately two week intervals up to 6 times per season. Apply no more than 3 quarts per season.</p>
<p>FERTIGATION RATES: The number and frequency of applications determine application rates. Apply 1-1.5 qt/ac/application, up to 3 times per season. Always apply toward the end of an irrigation cycle.</p>
<p>CROPS: Forage Crops: Alfalfa, Clover, Hay Crops; Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane; Legumes: Dry bean, Lentil, Peanut, Pea, Soybean; Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip; Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach; Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato; Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry; Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince; Other Fruit: Citrus (all), Fig, Kiwi, Melons; Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut; Grapes: Currant, Raisin, Table, Wine; Other Crops: Hemp, Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower</p>

WARNING

Causes eye irritation.
 Contains 2-3% of ingredients of unknown toxicity (inhalation and dermal).
 Wash hands and skin thoroughly after handling.
 If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
 If eye irritation persists: Get medical advice/attention.
 Dispose of contents/container in accordance with local, regional, national and international regulations.
 The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

Dispose of contents and container in accordance with local, regional, national and international regulations.

ATTENTION: The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

KEEP OUT OF REACH OF CHILDREN.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg is not liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.



Net Contents:

**Net Weight/
Gallon:**

Net Weight:

Batch No.:

Rev: 090920US-CA

NutriAg Ltd.

39 Gail Grove, Toronto, ON, Canada, M9M 1M5. Tel: (416) 636-1555

nutriag.com



INNOVATORS IN THE FIELD