



**ALIETTE® WDG brand Fungicide**  
**For Aerial Application to Strawberries**  
**EPA Reg. No. 264-516**  
**EPA SLN. No. CA-970008**

**Aventis CropScience USA LP**  
**P.O. Box 12014, 2 T.W. Alexander Drive**  
**Research Triangle Park, NC 27709**  
**(919) 549-2000**

## ***24(c) Supplemental Label***

### **For Use and Distribution Only in the State of California**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

<b>Crop/Site:</b>	Strawberry / California
<b>Target Pest:</b>	Red Stele ( <i>Phytophthora fragariae</i> ); Leather Rot ( <i>Phytophthora cactorum</i> )
<b>Rate:</b>	2.5 - 5.0 lbs product per acre
<b>Dilution Rate:</b>	Minimum of 10 gallons per acre
<b>Method of Application:</b>	Aerial
<b>Frequency and Timing of Application:</b>	Do not apply more than 30 lbs per season. For Red Stele begin applications 14-21 days after planting and continue on a 30-60 day interval as long as conditions favor disease development. For Leather Rot begin applications between 10% bloom and early fruit set and continue on a 7-14 day interval as long as conditions are favorable for disease development. The shorter time intervals and higher rates should be used when disease pressure is high.
<b>Field Reentry Safety Interval:</b>	Refer to the EPA registered label for ALIETTE® WDG Fungicide (EPA No. 264-516) for safety and re-entry information.

#### **Other Requirements:**

**Note:** Aventis CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG or if unbuffered ALIETTE® WDG is applied to foliage with copper residues. Do not tank-mix with copper fungicides.

When applied prior to or after copper fungicides, the pH of ALIETTE® WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG).

Adjuvants which enhance pesticide penetration may cause phytotoxicity when mixed with ALIETTE® WDG.

Mixing ALIETTE® WDG with surfactants or foliar fertilizers is not recommended. Strawberry sensitivity to a particular combination should be determined by spraying a few plants. Evaluate 3-7 days later for adverse effects.

ALIETTE is a registered trademark of the Aventis Group.