



**Bayer CropScience LP**  
**P.O. Box 12014**  
**2 T.W. Alexander Drive**  
**Research Triangle Park, North Carolina 27709**  
**1-866-99BAYER (1-866-992-2937)**

**ALIETTE® WDG Brand Fungicide**

**For Suppression of Citrus greening / HLB in Bearing and Non-Bearing Citrus**

**EPA Reg. No. 264-516**

***FIFRA 2(ee) Recommendation***

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF FLORIDA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

**CITRUS (BEARING AND NON-BEARING)**

DISEASE	APPLICATION METHOD	RATE product	APPLICATION PROGRAM
<b>Citrus greening / HLB</b> (Huanglongbing) (Candidatus Liberibacter asiaticus)	Foliar	2.5 - 5.0 lbs/acre	Pre-Bloom to May
			Mid-Aug to December

**RESTRICTIONS AND LIMITATIONS:**

- Any combination of labeled application methods can be used, however, do not exceed a total of four applications or 20 lbs/Acre per year.
- Do not exceed 100 GPA for foliar applications on non-bearing citrus and 500 GPA for foliar applications on bearing citrus.
- Do not graze livestock in treated groves.
- Do not feed forage from treated groves.

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG Brand Fungicide or if unbuffered ALIETTE® WDG Brand Fungicide is applied to foliage with copper residues. Do not tank-mix with copper compounds. When applied prior to or after copper compounds, the pH of ALIETTE® WDG Brand Fungicide should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG Brand Fungicide) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG Brand Fungicide). Adjuvants which enhance pesticide penetration may cause phytotoxicity when mixed with ALIETTE® WDG Brand Fungicide. Mixing ALIETTE® WDG Brand Fungicide with surfactants or foliar fertilizers is not recommended. Citrus sensitivity to a particular combination should be determined by spraying a small area of foliage and fruit. Evaluate 3-7 days later for adverse effects

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*