



FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN ALABAMA

Application in citrus groves to trees established less than one year

Alion[®] Herbicide

EPA Reg. No. 264-1106

SLN. No. AL210001

THIS LABEL IS VALID UNTIL October 5, 2026 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

APPLICATION DIRECTIONS FOR USE IN CITRUS

Only apply Alion Herbicide in citrus groves where the soil has completely settled around citrus trees and there are no open channels or depressions in the soil that would allow the product to move into the root zone through open channels.

Citrus Crops: Crop group 10-10 including Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; clementine; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and or hybrids of these

Dose Rate Chart for Citrus

| Soil Texture | Alion Herbicide (fl oz product / broadcast acre) |
|--|---|
| Any soil except those that contain 20% or greater gravel content | 5 .0 to 6 .5 fl oz/A (0 .065 to 0 .085 lb ai/A) |

USE RESTRICTIONS

- **Do not apply** more than 10 .3 fl oz product/A (0 .134 lb ai/A) per year or in a 12 month period.
- When making more than one application per year, allow a minimum of 90 days between applications.

USE PRECAUTIONS

Use in Established Groves:

Only apply Alion Herbicide in groves where the trees have been established for a minimum of one year after transplanting.

Use in Recently Planted Citrus:

Alion Herbicide may be used in groves planted a minimum of one month provided the following condition exists:

- 1) The transplanted trees were potted plants (such as citripots) and not bare-rooted.
- 2) The trunks are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- 3) The trees are actively growing and exhibiting good health and vigor.

Avoid direct or indirect spray contact with crop foliage, green bark, roots, or fruit as it may cause localized crop injury or death. Only the trunks of trees transplanted more than one year may be sprayed with Alion Herbicide if the trunk is callused, mature brown bark. Contact of Alion Herbicide with tissues other than mature brown bark can result in serious damage or plant death.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience, LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167**