

**FIFRA Section 24(c) Special Local Need Label****FOR DISTRIBUTION AND USE ONLY WITHIN NEW JERSEY****Alion[®] Herbicide**

For the use of Alion Herbicide on Dormant Highbush Blueberry in Sandy Soils containing greater than 1% Organic Matter.

EPA Reg. No. 264-1106

SLN. No. NJ-230001

THIS LABEL IS VALID UNTIL DECEMBER 31, 2026 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ACTIVE INGREDIENT:

Indaziflam* 19.05%

OTHER INGREDIENTS: 80.95%**TOTAL:** **100.00%**

Contains 1.67 pounds of indaziflam per gallon.

*(CAS No: 730979-19-8)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

APPLICATION DIRECTIONS FOR USE ON BLUEBERRY (Highbush) AND cultivars, varieties, and/or hybrids of these including rabbiteye blueberries, except blueberry (lowbush).

Only use Alion Herbicide in established plantings at least one year after the bushes have been planted and exhibiting normal growth and good vigor. Only apply Alion Herbicide to blueberries and bushberries where the soil has completely settled around the bushes and there are no exposed roots, open channels or depressions in the soil that would allow the product to move into the root zone or injury may occur.

Dose Rate Chart for Blueberry (Highbush)

Soil Texture	Soil Percent Organic Matter Content (%)	Rate Per Application (fl oz/A)	Max Rate Per Year (fl oz/A)	Minimum Plant Age
Sand	<1	DO NOT USE		1 year
	≥ 1	5.0 (0.065 lb ai/A)	10.0 (0.13 lb ai/A)	

USE RESTRICTIONS

- **Do not apply** more than a total of 10.0 fl oz product/A (0.13 lb ai/A) per year on sand containing ≥1 % organic matter content in a 12-month period when used in any highbush blueberry.
- **Do not use** in Blueberry (Highbush) crops grown in sand containing <1 % organic matter content.
- **Do not use** on soils with 20% or more gravel content.

- **Do not allow** spray to contact green stems, foliage, flowers, or berries or unacceptable injury may occur.
- **Only apply** Alion Herbicide to soil as a dormant application in late fall through early spring before bud swell.
- Apply Alion Herbicide as a directed application to the soil beneath the bushes.
- When making more than one application per year, allow a minimum of 90 days between applications.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call **1-866-99BAYER (1-866-992-2937)**.

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).



Bayer CropScience

**Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)**