



GUARANTEED ANALYSIS	
Total Nitrogen (N)	6.0%
5.5% Ammoniacal Nitrogen	
0.20% Nitrate Nitrogen	
0.30% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	20.0%
Zinc (Zn)	0.20%
0.20% Water Soluble Zinc (Zn)	
	monium Nitrate, Phosphoric Acid, and Zinc Gluconate
Derived From: Anhydrous Ammonia, Urea-Ami ALSO CONTAINS NON-PLANT FOOD INGRE Soil Amending Ingredients:	DIENTS
Derived From: Anhydrous Ammonia, Urea-Ami ALSO CONTAINS NON-PLANT FOOD INGRE Soil Amending Ingredients: 5.50%	DIENTSOrganic Matter (derived from leonardite)
Derived From: Anhydrous Ammonia, Urea-Ama ALSO CONTAINS NON-PLANT FOOD INGRE	DIENTSOrganic Matter (derived from leonardite)
Derived From: Anhydrous Ammonia, Urea-Ami ALSO CONTAINS NON-PLANT FOOD INGRE Soil Amending Ingredients: 5.50%	DIENTSOrganic Matter (derived from leonardite)
Derived From: Anhydrous Ammonia, Urea-Am ALSO CONTAINS NON-PLANT FOOD INGRE Soil Amending Ingredients: 5.50%	DIENTSOrganic Matter (derived from leonardite)
Derived From: Anhydrous Ammonia, Urea-Ami ALSO CONTAINS NON-PLANT FOOD INGRE Soil Amending Ingredients: 5.50%	Organic Matter (derived from leonardite)

KEEP OUT OF REACH OF CHILDREN

WARNING

Avoid breathing mists. Harmful if swallowed. Avoid contact with skin, eyes or clothing. FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.

Product is guaranteed effective up to

F1548

DIRECTIONS FOR USE

CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING
D1) Root and Tuber Vegetables including Carrot, binseng, Horseradish, Parsley (turnip-rooted), Potato, Radish, Sugar Beet, Sweet Potato, Turnip	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be ber eficial.
		Foliar: (minimum of 5 gpa) is an effective foliar nutrient when mid or late seaso phosphate levels fall below optimum values. Repeat applications may be beneficial
03) Bulb Vegetables including Garlic, Leek, Onion	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be ber eficial.
		Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seaso phosphate levels fall below optimum values. Repeat applications may be beneficial
(04) Leafy Vegetables including Celery, Head and Leaf Lettuce and Spinach	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be ber eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seaso phosphate levels fall below optimum values. Repeat applications may be beneficial.
(05) Brassica Vegetables including Broccoli, Cabbage, Cauliflower, Kale, and Mustard greens	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Follar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seaso phosphate levels fall below optimum values. Repeat applications may be beneficial.
(06) Legume Vegetables (Succulent or Dried) including Bean all types), Peas and Soybeans	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be ber eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seaso phosphate levels fall below optimum values. Repeat applications may be beneficial.
(08) Fruiting Vegetables including Eggplant, Pepper and Tomato	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be ber eficial. Foliar: (minimum of 5 gpa) is an effective foliar nutrient when mid or late seasc phosphate levels fall below optimum values. Repeat applications may be beneficial.
09) Cucurbit Vegetables including Cantaloupe, Cucumber, Honeydew, Muskmelon, Squash summer and winter) and Watermelon	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Foliar: (minimum of 5 gpa) is an effective toliar nutrient when mid or late sease phosphate levels fall below polimum values. Repeat applications may be beneficial.
(10) Citrus Fruit including Grapefruit, Lemon, Lime, Sweet Orange and Tangerine	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soll: May be placed surface or subsurface band. Continue irrigation to flush line and maximize nutrient distribution. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa) is an effective foilar nutrient when mid or late seasc phosphate levels fall below optimum values. Repeat applications may be beneficial.
(11) Pome Fruits including Apple and Pear	Soil: 1-20 gal/ac Foliar: 2-16 qt/ac	Soil: May be placed surface or subsurface band. Continue irrigation to flush line and maximize nutrient distribution. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa) is an effective foliar nutrient when mid or late sease phosphate levels fall below optimum values. Repeat applications may be beneficial.
(12) Stone Fruits including Apricot, Cherry, Peach and Plumcot	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Continue irrigation to flush line and maximize nutrient distribution. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa), is an effective foliar nutrient when mid or late sease phosphate levels fall below optimum values. Repeat applications may be beneficial.
(13) Berry and Small Fruit including Blackberry, Blueberry, Caneberry, Kiwi, Raspberry and Grape	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Continue irrigation to flush line and maximize nutrient distribution. Repeat applications may be beneficial. Foliar: prinimum of 5 gpa), is an effective foliar nutrient when mid or late seasc phosphate levels fall below optimum values. Repeat applications may be beneficial.
14) Tree Nuts including Almonds, Cashews and Pecans	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Continue irrigation to flush line and maximize nutrient distribution. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa), is an effective foliar nutrient when mid or late sease phosphate levels fall below optimum values. Repeat applications may be beneficial.
15) Grain, Cereal	Soil: 1-20 gal/ac Foliar: 2-16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seas phosphate levels fall below optimum values. Repeat applications may be beneficial.
16) Forage, Fodder and Straw of Cereal Grains	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seass phosphate levels fall below optimum values. Repeat applications may be beneficial.
17) Grass Forage, Fodder and Hay	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seass phosphate levels fall below optimum values. Repeat applications may be beneficial.
18) Nongrass Animal Feeds including Alfalfa, Clover, Hay and Vetch	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be be eficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late seass phosphate levels fall below optimum values. Repeat applications may be benefici
10) Harba and Chicae including Pacil Dill Mustard	Cail: 1 20 gal/aa	Call. May be placed surface or subsurface hand. Depost applications may be be

Directions for use cont'd.			
CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING	
(22) Stalk, Stem and leaf petiole vegetables, Asparagus, bamboo shoots and celery	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late season phosphate levels fall below optimum values. Repeat applications may be beneficial.	
Cotton, Peanut, Sugarcane, Tobacco	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late season phosphate levels fall below optimum values. Repeat applications may be beneficial.	
Turf and Ornamental	Soil: 1-20 gal/ac Foliar: 2- 16 qt/ac	Soil: May be placed surface or subsurface band. Repeat applications may be beneficial. Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late season phosphate levels fall below optimum values. Repeat applications may be beneficial.	

Soil Application Rates:
Altura NST maybe applied pre-plant, as a starter, side dress, fertigated, top dressed or banded on the soil. Dilute water (If necessary) to ensure uniform distribution.

retrigation Application Rates:
Continue irrigation after application of Altura NST for a minimum of 2 hours to flush lines and to maximize nutrient distribution with drip and micro sprinkler use. Repeat applications may be beneficial.

Application Through Drip Irrigation – inject after irrigation filter.

Transplant Water: Use 3-10 gallons per acre, not to exceed 3% solution of Altura NST to water.

Apply Altura NST in enough water for adequate coverage (5 gpa minimum). A maximum concentration of (1:5 Altura NST to water) should be used for foliar applications. May be applied through center pivot irrigation systems. Repeat applications may be beneficial. STORAGE AND DISPOSAL

STORAGE: Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal. DISPOSAL: Do not reuse container. Completely empty container into application equipment. Thoroughly rinse container, adding rinse water to the spray tank.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300. CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR

INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY. Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-lpi.com/

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible

Altura NST is a registered trademark of Loveland Products, Inc.

WARNING

This product can expose you to chemicals including cadmium, which are known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov.

2 years after date stamped on container.

Density: 10.8 lbs/gallon (1,295 grams/liter) @ 68°F

Net Weight: 265 gallons (1003.1 liters), 2,862 Lbs (1,298 kg)

Product Code: 1000992346

Density: 10.8 lbs/gallon (1,295 grams/liter) @ 68°F

(19) Herbs and Spices including Basil, Dill, Mustard and Sage Foliar: 2- 16 qt/ac

(20) Oil Seed Crops including Canola, Flax and

gallons (**Net Contents:** liters) **Net Weight:** lbs (Product Code: 1000992348

AlturaNST D3021

Soil: May be placed surface or subsurface band. Repeat applications may be ber Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late season phosphate levels fall below optimum values. Repeat applications may be beneficial.

Soil: May be placed surface or subsurface band. Repeat applications may be ben-Foliar: (minimum of 5 gpa).is an effective foliar nutrient when mid or late season phosphate levels fall below optimum values. Repeat applications may be beneficial.