

GROUP 1 FUNGICIDE

PULL HERE TO OPEN ►



syngenta®

Fungicide

Active Ingredient:

Thiabendazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*CAS No.148-79-8

Alumni is a suspension concentrate containing 4.1 pounds of Thiabendazole per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

[See additional precautionary statements and directions for use inside booklet.]

EPA Reg. No. 100-889

EPA Est. 39578-TX-1

Product of India

Formulated in the USA

SCP 889C-L1C 0715

4057483

2.5 gallons
Net Contents

®

FIRST AID	
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS	
Hazards to Humans and Domestic Animals	CAUTION
Harmful if inhaled. Avoid breathing vapor or spray mist. Remove and wash contaminated clothing before reuse.	
Personal Protective Equipment (PPE)	
Mixers, loaders, applicators and other handlers must wear:	
<ul style="list-style-type: none"> • Long-sleeved shirt and long pants • Shoes plus socks 	
User Safety Requirements	
Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.	
User Safety Recommendations	
Users should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticides gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	
Environmental Hazards	
This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.	

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

Alumni is a systemic fungicide which aids in the control of many important plant diseases. If treatment is not effective following use of Alumni as recommended, a tolerant strain of fungus may be present. Consideration should be given to prompt use of other suitable non-benzimidazole fungicides. Alumni is not for use in formulating agricultural products.

RESISTANCE MANAGEMENT

GROUP 1 FUNGICIDE

This product contains Thiabendazole, a Group 1 fungicide. Thiabendazole belongs to the methyl-benzimidazole carbamate class of chemistry which disrupts β -tubuline assembly in mitosis.

Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include alternation and/or tank-mixing with products having different modes of action or limiting the total number of applications per season.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local extension specialist or certified crop advisor

Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>

CROP USE DIRECTIONS

GENERAL – SHAKE OR STIR WELL BEFORE USE.

Clean equipment before using Alumni. **When Alumni is dispersed in water, the resulting suspension must be constantly agitated.** Avoid foaming.

Use the higher specified rates under conditions of heavier disease pressure.

Compatibility information for Alumni with other chemicals is limited. Consult your supplier or agricultural experiment station for specific recommendations.

Citrus (One Application Only) – Blue mold (*Penicillium digitatum*), Green mold (*Penicillium italicum*), and stem end rot (*Diplodia natalensis*): Dip, flood, or spray harvested fruit with a suspension of Alumni in water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion. Rate will vary according to application method; please refer to the following for specific rates for citrus blue mold control.

Citrus Bin Dips, Floods and Drenches

Use 26 fl oz of Alumni in 100 gal of water to provide 1,000 ppm Thiabendazole. Citrus must be drenched for three (3) minutes. Use one (1) gal of liquid per 1,500-2,000 lb citrus or 100 gal of liquid for about 30 bins. Avoid excessive foaming, use antifoam if needed.

Citrus Sprays

Use 52-91 fl oz Alumni in 100 gal of water, wax/oil emulsions, or an aqueous dilution of wax/oil emulsion to provide 2,000-3,500 ppm of Thiabendazole. Keep the mixture under constant agitation. Apply at a rate of one gal per 8,000-10,000 lb citrus fruit.

For sporulation control, apply 130 fl oz of Alumni in 100 gal of water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion to provide 5,000 ppm. Apply at a rate of one gal per 8,000 – 10,000 lb citrus fruit.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store Alumni in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

STORAGE AND DISPOSAL (continued)

Container Handling [greater than 5 gallons – mini-bulk]



Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons - bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Alumni®, the ALLIANCE FRAME 
the SYNGENTA Logo and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

©2015 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 889C-L1C 0715
4057483

GROUP **1** FUNGICIDE



Fungicide

Active Ingredient:	
Thiabendazole*:	42.3%
Other Ingredients:	57.7%
Total:	100.0%

*CAS No.148-79-8

Alumni is a suspension concentrate containing 4.1 pounds of Thiabendazole per gallon.

[See additional precautionary statements and directions for use inside booklet.]

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-889
EPA Est. 39578-TX-1

Alumni® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

©2015 Syngenta

Manufactured by:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 889C-L1C 0715
4057483

KEEP OUT OF REACH OF CHILDREN. CAUTION

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call **1-800-888-8372**.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store Alumni in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

2.5 gallons
Net Contents

syngenta®

