



DIFENOCONAZOLE TEBUCONAZOLE	GROUP	3	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE

APPRISE™

Seed Treatment

FOR USE ON BARLEY, OATS, TRITICALE, AND WHEAT FOR PROTECTION OF SEEDLINGS AGAINST LISTED SOILBORNE DISEASES

ACTIVE INGREDIENTS:

Difenoconazole: 3.57%
 Metalaxyl: 1.84%
 Tebuconazole: 0.39%

OTHER INGREDIENTS: 94.20%
TOTAL: 100.00%

Contains:

0.31 pound Difenoconazole per gallon (37.2 grams per liter)
 0.16 pound Metalaxyl per gallon (19.2 grams per liter)
 0.034 pound Tebuconazole per gallon (4.1 grams per liter)

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS,
 AND DIRECTIONS FOR USE

For Chemical Spill, Leak, Fire,
 or Exposure, Call CHEMTREC
 (800) 424-9300

For Medical Emergencies Only,
 Call (877) 325-1840

EPA Reg. No. 55146-160-48013
 EPA Est. No. 228-IL-002

Distributed by:
 CHS Inc.
 5500 Cenex Dr.,
 Inver Grove Heights, MN 55077

Net Contents
2.5 Gal.
(9.46 L)
Nonrefillable Container

15366000



FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride (PVC), Viton).

Note: Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT place product near or allow product to come into contact with strong oxidizing substances (such as potassium permanganate).

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to mammals, fish, and aquatic invertebrates. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from treated areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA. Cover or incorporate spilled treated seeds.





Groundwater Advisory Statement: Metalaxyl and Tebuconazole are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (RE) of 24 hours**.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinylchloride (PVC) or Viton.
- Shoes plus socks.

USE RESTRICTIONS

- **DO NOT** use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.
- Only for use in commercial seed treatment establishments in liquid or slurry treaters. Not allowed for "on-farm" use.
- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs. Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- **DO NOT** graze or feed livestock on green barley forage until 31 days after planting.
- **DO NOT** graze or feed livestock on green wheat, oats, or triticale for 55 days after planting.
- **DO NOT** use the same auger for treating seed that will handle commercial or feed grains.
- If excess treated seed remains after planting, dispose of it by burying away from bodies of water.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and, (2) no measurable residues remain in ethanol by-products that are used in agronomic practice.

ROTATIONAL CROPS

DO NOT plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.

RESISTANCE MANAGEMENT RECOMMENDATIONS

CHS Apprise seed treatment contains difenoconazole and tebuconazole, Group 3 fungicides and; metalaxyl, a Group 4 fungicide. Difenoconazole and tebuconazole belong to the DMI class of chemistry. Metalaxyl belongs to the phenylamide class of chemistry.

Fungal populations may contain individuals naturally resistant to active ingredients in Salient TMI and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as an alternation with fungicides of a different mode of action and/or tank mixes established for the crop and use area.





Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Nufarm encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact Nufarm Americas Inc representatives at 1-800-345-3330
- Contact your local extension specialist or certified crop advisor.
- Visit the Fungicide Resistance Action Committee (FRAC) at <http://www.frac.info>

PRODUCT INFORMATION

Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application to seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates for the crop to be treated with this product.

The following table lists the maximum recommended total slurry volumes by crop.

Crop	Maximum Fl. Oz. Slurry / 100 lbs. of Seed
Wheat, Oats*, Triticale, and Barley	16

* Not approved for use in California

SEED TREATMENT

This product is a mixture of the fungicides Difenoconazole, Tebuconazole, and Metalaxyl. When used on listed crops, this product aids in the control or suppression of the following seed, seedling and soilborne diseases on seed and young seedlings of Barley, Oats, Triticale, and Wheat.

CROP	DISEASE CONTROL	RATE
Barley Oats* Triticale Wheat	Control of seed rots such as <i>Penicillium</i> and <i>Aspergillus</i> and seed-borne <i>Fusarium</i> Scab and <i>Septoria</i> . Control of Dwarf bunt, Karnal bunt*, Stinking smut (Common bunt), Flag smut, Loose smut, Covered smut of barley. Control of early season Barley stripe, <i>Pythium</i> damping-off, <i>Septoria</i> disease complex, <i>Rhizoctonia</i> root rot, Common root rot (<i>Cochliobolus</i> spp.), <i>Fusarium</i> foot rot. Early season suppression of powdery mildew, wheat leaf rust, and <i>Septoria</i> leaf blotch.	5.0 – 7.5 fluid ounces per 100 pounds of seed
<p>APPLICATION DIRECTIONS: Apply prior to planting as a slurry treatment in sufficient fluid for adequate coverage. This product is to be used in liquid or slurry treaters only. Mix thoroughly before use or use entire container at one time. Ensure thorough coverage.</p> <p>TANK MIXES: All tank mixes should be pre-tested to determine physical compatibility between formulations. Follow the most restrictive precautions and limitations on the labeling of all products used in mixtures. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>*Not approved for use in California</p>		

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with CHS Apprise seed treatment (difenoconazole, metalaxyl, and tebuconazole).
- **DO NOT** use for feed, food, or oil purpose.

THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH CHS APPRISE SEED TREATMENT (difenoconazole, metalaxyl, and tebuconazole):

- **Groundwater Advisory:** Metalaxyl and Tebuconazole are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Store treated seed away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves when opening this bag or handling (e.g. loading, pouring) treated seed.





- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** graze or feed livestock on green barley forage until 31 days after planting.
- **DO NOT** graze or feed livestock on green wheat, oats, or triticale for 55 days after planting.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **DO NOT** plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Store treated seed in a cool, dry location.
- **DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place, **DO NOT** store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less

Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons

Nonrefillable container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Refillable containers larger than 5 gallons

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. For final disposal, offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.





DISCLAIMER OF WARRANTIES

Information, descriptions, and illustrations in brochures, leaflets, or other marketing materials are based as closely as possible on experiences in tests and in practice and do not constitute quality claims, representations, and/or warranties. CHS Inc. and seller do not warrant, guarantee, or imply similar results. Tests have not been conducted under all possible conditions or agronomic practices, and CHS Inc. and seller do not warrant or represent that the product is usable under all possible use conditions. The purchaser and/or user must determine whether the product is suitable for the intended purpose and/or can be used in the local conditions with the user's specific agronomic practices.

CHS Inc. and seller, as their sole and exclusive warranty, state that when the product was shipped from a CHS Inc. facility, the product conformed to the descriptions on the product label, within tolerances allowed by law. CHS INC. AND SELLER MAKE NO OTHER WARRANTIES OR REPRESENTATIONS REGARDING THE PRODUCT, AND CHS INC. AND SELLER DISCLAIM ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY AND WARRANTIES FOR A PARTICULAR PURPOSE. THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL OF THE PRODUCT CONTAINER.

No person or agent on behalf of CHS Inc. or seller has any authority to make any warranties or representations different from, or in addition to, the sole and exclusive warranty stated in these Terms and Conditions of Sale.

LIMITATION OF LIABILITY, ALLOCATION OF RISK, AND LIMITATION OF REMEDIES

No Guarantee of Yield or Crop Performance, and Recovery Limited to Purchase Price

Crop yield and quality are dependent on many factors beyond CHS Inc.'s and seller's control. These factors include, but are not limited to, weather, agronomic practices, manner of use, soil conditions, presence of other materials, disease, pests, handling, and storage. CHS Inc. and seller do not guarantee or make any warranties regarding yields or crop performance. CHS Inc. and seller do not represent that the product is free from latent defects unknown to CHS Inc. and seller at the time of sale. These are all inherent risks in growing crops, and all risks of nonperformance, reduced performance, and/or crop damage are assumed by the purchaser and the user.

PURCHASER OR USER MAY NOT RECOVER FROM CHS INC. OR SELLER ANY AMOUNT FOR INCIDENTAL, INDIRECT, CONSEQUENTIAL OR PUNITIVE DAMAGES, INCLUDING LOSS OF PROFIT, LOSS OF YIELD, AND AMOUNTS EXPENDED IN USING THE PRODUCT, OR FOR GROWING, HARVESTING OR MARKETING THE PRODUCE FOR WHICH THE PRODUCT WAS PURCHASED. THIS LIMITATION OF LIABILITY IS APPLICABLE TO ANY CLAIM, WHETHER THE LEGAL THEORY FORMING THE BASIS OF SUCH CLAIM INVOLVES CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE.

PURCHASER'S AND/OR USER'S SOLE AND EXCLUSIVE REMEDY, AND CHS INC.'S AND SELLER'S SOLE LIABILITY FOR LOSS OR DAMAGE ARISING FROM THE PURCHASE AND/OR USE OF PRODUCT FROM CHS INC. IS A REFUND OF PURCHASE PRICE OF THE PRODUCT. Purchaser and user acknowledge and agree that limiting any potential recovery to the purchase price of the product is an appropriate allocation of risk between CHS Inc., seller, and the purchaser and/or user given the risks and uncertainties inherent in the agricultural business outside of CHS Inc.'s and seller's control such as weather, agronomic practices, planting, cultivating, harvesting, and marketing decisions. Purchaser and user acknowledge and agree that limiting any potential recovery to the purchase price of the product is reflected in the purchase price of the product, and without this limitation the purchase price of the product would be significantly higher.

PURCHASER AND USER ACKNOWLEDGE AND AGREE THAT IF CHS INC. OR SELLER REFUND THE PURCHASE PRICE OF THE PRODUCT, THE LIMITATION WILL NOT HAVE FAILED OF ITS ESSENTIAL PURPOSE.

These warranty disclaimers, allocation of risk, and limitations are reflected in the price of the product. If the purchaser or user would like a different warranty, allocations of risk, or limitations, which will substantially increase the price of the product, purchaser or user may contact CHS Inc. to discuss different terms and a different product price. If no agreement is reached, then any unopened product can be returned pursuant to the above terms.

PROMPT NOTICE OF CLAIM:

CHS Inc. must have prompt notice of any claim arising from the use of the product so that a timely inspection of the alleged affected product and/or crop can be made. Purchaser or user shall notify CHS Inc. immediately, but in no case later than thirty (30) days after any defect or other basis of such claim is discovered or should have been discovered. Failure of the purchaser or user to provide notice to CHS Inc. within thirty (30) days will constitute a waiver, and after such time all claims will be barred.

NO AMENDMENT WITHOUT SIGNED WRITING:

The Terms and Conditions of Sale may not be altered or amended in any way except in writing, and signed by an officer of CHS Inc.

RV060718

CHS Apprise™ is a trademark of CHS Inc.

All other trademarks are the property of their respective owners.





NOTES





NOTES





DIFENOCONAZOLE	GROUP	3	FUNGICIDE
TEBUCONAZOLE			
METALAXYL	GROUP	4	FUNGICIDE



APPRISE™

Seed Treatment

FOR USE ON BARLEY, OATS, TRITICALE, AND WHEAT FOR PROTECTION OF SEEDLINGS AGAINST LISTED SOILBORNE DISEASES

ACTIVE INGREDIENTS:

Difenoconazole:	3.57%
Metalaxyl:	1.84%
Tebuconazole:	0.39%
OTHER INGREDIENTS:	94.20%
TOTAL:	100.00%

Contains:

- 0.31 pound Difenoconazole per gallon (37.2 grams per liter)
- 0.16 pound Metalaxyl per gallon (19.2 grams per liter)
- 0.034 pound Tebuconazole per gallon (4.1 grams per liter)

PULL HERE TO OPEN ↑

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ATTACHED BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, AND DIRECTIONS FOR USE

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. For Medical Emergencies Only, Call (877) 325-1840.

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by the poison center or doctor. DO NOT give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in a cool place, **DO NOT** store in direct sunlight. Protect from freezing temperatures.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

(continued)

STORAGE AND DISPOSAL (continued)

CONTAINER HANDLING: NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less

Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons

Nonrefillable container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Refillable containers larger than 5 gallons

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. For final disposal, offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Distributed by:
CHS Inc.

5500 Cenex Dr.,
Inver Grove Heights, MN 55077

EPA Reg. No. 55146-160-48013
EPA Est. No. 228-IL-002

RV060718

