

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1476A-S1 1217**

<b>BENZOVINDIFLUPYR</b>	<b>GROUP</b>	<b>7</b>	<b>FUNGICIDE</b>
<b>DIFENOCONAZOLE</b>	<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>

### **Aprovia® Top Fungicide**

**This supplemental label expires on 12/18/2020 and must not be used or distributed after this date.**

Active Ingredients:	
Difenoconazole* .....	10.95%
Benzovindiflupyr** .....	7.30%
<hr/>	
Other Ingredients:	81.75%
<hr/>	
Total:	100.00%

\*CAS No. 119446-68-3

\*\*CAS No. 1072957-71-1

Aprovia Top Fungicide is formulated as an emulsifiable concentrate containing 0.97 lb ai of difenoconazole active ingredient and 0.65 lb ai of benzovindiflupyr active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1476

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Aprovia Top Fungicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**



## DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate Fl Oz Product/A	Remarks
<b>Bulb Crop Group 3-07</b>	Cladosporium leaf blotch ( <i>C. allii</i> )	10.5*	Applications should begin prior to disease development and continue throughout the season on a 7-14 day schedule.
<b>Dry Bulb Group 3-07A</b>	Powdery Mildew ( <i>Leveillula taurica</i> )		
Onion	Purple Blotch ( <i>Alternaria porri</i> )		No more than two applications of Aprovia Top Fungicide may be applied on a 7-day interval. All other applications must be applied no closer than a 14-day interval. For resistance management, do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide.
Garlic	Rust ( <i>Puccinia allii</i> )		
Shallot (see others below)	Stemphyllium leaf blight and stalk rot ( <i>S. vesicarium</i> )		
<b>Onions, green Group 3-07B</b>	Suppression:		The addition of a spreading/penetrating type adjuvant such as organo-silicon blends with either non-ionic surfactants (NIS) or vegetable based crop oils (COC); or vegetable based COC (not mineral); or NIS with at least 90% concentration is recommended.
Chives	Rhizoctonia seedling disease ( <i>R. solani</i> )		
Leeks			If disease pressure is high, use the shortest interval.
Green			

\*10.5 fl oz product/A is equivalent to 0.080 lb ai difenoconazole and 0.053 lb ai benzovindiflupyr.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Aprovia Top Fungicide can be applied by ground, air, or chemigation. Use a minimum of 10 gal/A for ground applications is recommended. Use a minimum of 5 gal/A for aerial applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

**Complete List of Bulb Crop (dry bulb) Crop Group 3-07A:** Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these

**Complete List of Green Onion Crop Group 3-07B:** chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these

### Specific Use Restrictions:

- 1) Do not apply more than 54 fl oz/A/year of Aprovia Top Fungicide (equivalent to 0.412 lb ai difenoconazole and 0.272 lb ai benzovindiflupyr).
- 2) Do not exceed 4 applications per year.
- 3) Do not exceed 13.5 fl oz product per application.
- 4) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products for dry bulb onions or more than 21 fl oz/A/year (0.34 lb ai/A/year) for green onions.
- 5) Do not apply more than 0.27 lb ai/A/year of benzovindiflupyr-containing products.
- 6) Aerial application is prohibited in New York State.
- 7) Do not apply within 7 days of harvest (7 day PHI).

©2018 Syngenta

Aprovia® Top and the Syngenta logo are trademarks of a Syngenta Group  
Company