

## SUPPLEMENTAL LABELING



TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE

**EPA Reg. No. 71711-36**

### **Bushberry Subgroup (Crop Subgroup 13-07B)**

aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry; native currant; salal; sea buckthorn; cultivars, varieties, and/or hybrids of these

This supplemental labeling expires 03/31/2022 and must not be used or distributed after this date.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for APTA® Insecticide before applying. Use of APTA Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for APTA Insecticide. New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

### **APPLICATION RATE CHART FOR APTA INSECTICIDE**

Pest	Rate/Acre	Use Directions
Aphids Azalea bark scale Blueberry blossom weevil Blueberry gall midge Blueberry maggot Blueberry spanworm Cranberry fruitworm Cherry fruitworm Eriophyid mites Katydid Leafhoppers Lecanium scale Oblique-banded leafroller Plum curculio Powdery mildew Thrips, including citrus thrips Spotted wing drosophila*	27.0 fl oz/acre (0.28 lb ai/acre)	<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• <b>Do not apply by air on bushberry.</b></li> <li>• Apply by ground <b>only</b>, using a minimum of 20 gallons of water per acre.</li> <li>• Do not apply more than 81.0 fl oz (0.85 lb ai) per acre per year.</li> <li>• Do not make more than 3 applications per year.</li> <li>• Allow at least 14 days between applications.</li> <li>• Preharvest Interval (PHI): 3 days</li> </ul> <b>USE RECOMMENDATIONS - DISEASES</b> <ul style="list-style-type: none"> <li>• Begin applications prior to onset of disease.</li> <li>• Use of an agriculturally approved nonionic surfactant at 0.25% v/v may improve disease control.</li> <li>• If weather conditions remain conducive to disease development, apply another registered fungicide product with a different mode of action 7 to 10 days later.</li> <li>• Consult local extension recommendations or your agricultural consultant for information specific to your area.</li> </ul>
*suppression only		

©2019 Nichino America, Inc. Apta and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO**  
**AMERICA**

Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700