SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



EPA Reg. No. 71711-36

Pome Fruit (Crop Group 11-10)

apple; azarole; crabapple; loquat; mayhaw; Hook. & Arn., and *C. rufula* Sarg; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for APTA insecticide before applying. Use of APTA insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for APTA insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART FOR APTA INSECTICIDE

Pest	Rate/Acre	Use Directions
Katydid Leafhoppers	14.0 to 21.0 fl oz/acre (0.15 to 0.22 lb ai/acre)	Apply by ground only, using a minimum of 100 gallons of water per
Aphids (excluding woolly apple aphid)	17.0 to 21.0 fl oz/acre (0.18 to 0.22 lb ai/acre)	 Do not apply by air on pome fruit. Do not apply more than 53.5 fl oz (0.56 lb ai) per acre per year. Do not make more than 2 applications
Apple maggot Leafrollers Mealybugs Pear psylla Pear rust mite Plum curculio Thrips	21.0 to 27.0 fl oz/acre (0.22 to 0.28 lb ai/acre)	 Allow at least 14 days between applications. Preharvest Interval (PHI): 14 days
Codling moth* Powdery mildew* *suppression only		

©2018 Nichino America, Inc. APTA and Nichino America logo are registered trademarks of Nichino America, Inc.