

## SUPPLEMENTAL LABELING

|             |       |     |             |
|-------------|-------|-----|-------------|
| TOLFENPYRAD | GROUP | 21A | INSECTICIDE |
|             | GROUP | 39  | FUNGICIDE   |



EPA Reg. No. 71711-36

### Citrus Fruit (Crop Group 10-10)

Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin orange, clementine); tangor; trifoliate orange; uni fruit; cultivars, varieties, and/or hybrids of these

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for APTA insecticide before applying. Use of APTA insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for APTA insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

### APPLICATION RATE CHART FOR APTA INSECTICIDE

| Pest   | Rate/Acre  | Use Directions  |
|--|--|---|
| Asian citrus psyllid<br>Katydid  | 14.0 to 27.0 fl oz/acre<br>(0.15 to 0.28 lb ai/acre) | <ul style="list-style-type: none"> <li>For <b>high air velocity, low volume or air curtain sprayers</b>, apply using a minimum of 30 gallons of water per acre</li> </ul> <b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li><b>Do not apply by air on citrus fruit.</b></li> <li>Apply by ground <b>only</b>, using a minimum of 100 gallons of water per acre.</li> <li>Do not use on nursery stock.</li> <li>Do not apply more than 27.0 fluid ounces (0.28 lb ai) per acre per growing season.</li> <li>Do not make more than 1 application per growing season.</li> <li>Do not make more than 2 applications per year.</li> <li>Preharvest Interval (PHI): 3 days</li> </ul> |
| Aphids   | 17.0 to 27.0 fl oz/acre<br>(0.18 to 0.28 lb ai/acre) |   |
| Citrus bud mite<br>Citrus mealybug<br>Citrus thrips<br>Soft scale insects, including<br>citricola scale, barnacle<br>scale, brown soft scale | 21.0 to 27.0 fl oz/acre<br>(0.22 to 0.28 lb ai/acre) |   |
| Citrus peelminer<br>Citrus red mite<br>Citrus rust mite (silver mite)<br>Mealybugs<br>Citrus orangedog*<br>Citrus weevil*<br>Cutworms*       | 24.0 to 27.0 fl oz/acre<br>(0.25 to 0.28 lb ai/acre) |   |
| *suppression only  |  |   |

©2018 Nichino America, Inc. APTA and Nichino America logo are registered trademark of Nichino America, Inc.

**NICHINO**  
**AMERICA**

Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700