



Below recommended rates in up to 100 gallons of water per acre.

CROP TYPE	PEST	RATE (Oz./Ac.)
Tomatoes, Potatoes Peppers	European red mite, Pacific spider mite, Strawberry spider mite, Twospotted spider mite.	32-48
Soybeans	Whitefly, (Bemisia spp.), Leafminers (Aphids)	32-48
Corn	European red mite, Pacific spider mite, Strawberry spider mite, Twospotted spider mite.	32-48
Cucurbits: Cantaloupe, Honeydew, Melons, Watermelon, Pumpkin, Zucchini	European red mite, Pacific spider mite, Strawberry spider mite, Twospotted spider mite, Whitefly, (Bemisia spp.), Leafminers (Aphids)	32-48
Turf and Ornamentals	Whitefly, (Bemisia spp.), Leafminers (Aphids)	32-48
Citrus	Asian citrus psyllid, Broad mite, Citrus bud mite, Citrus rust mite, Twospotted spider mite.	32-48
Alfalfa	Whitefly, (Bemisia spp.), Leafminers (Aphids)	32-48
Strawberries, Blueberries, Raspberries, Blackberries	Whitefly, (Bemisia spp.), Leafminers (Aphids)	32-48

**SHAKE WELL BEFORE USING. USE DILUTED SOLUTION IMMEDIATELY AFTER MIXING. DO NOT STORE DILUTED SOLUTION.**

**Application Instructions:** Do not apply more than two consecutive applications. To counter potential for resistance, rotate ARAMITE™ Organic Acaricide-Insecticide with a different product after two applications. Apply early in the morning or in late afternoon. Spray directly onto the insect or mite. Repeat applications at an interval of 7 days, depending on mite/insect pressure. To prevent product degradation, do not use spray water with a pH lower than 4.0 or higher than 8.0. Use pH correctors for pH's outside of these ranges.

**HAND HELD EQUIPMENT:** When using hand held equipment use 0.32-0.48/oz of ARAMITE® Organic Acaricide-Insecticide concentrate per gallon of spray water. Apply enough to cover the entire surface of infested tissues. Use higher rate for heavy infestations.

**PHYTOTOXICITY:** this product is not phototoxic and will not burn when used in accordance with label use instructions.