

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

Pressure – Use the lowest spray pressures recommended for the nozzle to produce the target spray volume and droplet size.

Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

Adjust nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

Boom Height - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift.

Boom-less Ground Applications: setting nozzles at the lowest effective height will help reduce the potential for spray drift.

Handheld Technology Applications: take precautions to minimize spray drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Wind

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Sensitive Areas

Apply this pesticide only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION INSTRUCTIONS

For optimum efficacy apply the spray solution in a manner that promotes maximum coverage, i.e. proper water volume, correct spray nozzles, optimum weed size, etc. Arroz 80 can be applied:

- By aircraft, with a total spray volume of 10 gallons or greater per acre
- By ground equipment with a total spray volume of 15 to 20 gallons or greater per acre

If spray volume is not sufficient, weed control can be compromised. If foliage canopy is heavy, use enough spray volume to reach and adequately cover weeds. Any factor that unfavorably affects weed coverage can result in compromised weed control. Buffer the application water if the pH is above 7.0 or below 6.0. Do not use turbid, high sediment or ditch water.

INSTRUCTIONS FOR USING WATER SOLUBLE PACKAGES DIRECTLY INTO SPRAY TANKS:

Water Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs. WSPs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

HANDLING INSTRUCTIONS

Follow these steps when handling pesticide products in WSPs:

- 1) Mix in spray tank only.
- 2) Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
- 3) Keep the WSP(s) in outer packaging until just before use.
- 4) Keep the WSP dry prior to adding to the spray tank.
- 5) Handle with dry gloves and according to the label instructions for PPE.
- 6) Keep WSP intact. Do not cut or puncture WSP.
- 7) Reseal the WSP outer package to protect any unused WSP(s)

MIXING INSTRUCTIONS

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

- 1) If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
- 2) Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
- 3) Stop adding water and stop any agitation.
- 4) Place intact/unopened WSP(s) into the tank.
- 5) Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
- 6) Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
- 7) Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
- 8) Stop agitation before tank lid is opened.

- 9) Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSP's have fully dissolved and the contents have been thoroughly mixed into the solution.
- 10) Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
- 11) Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
- 12) Use the spray solution when mixing is complete.
- 13) Maintain agitation of the diluted pesticide mix during transport and application.
- 14) It is unlawful to use any registered pesticide, including WSP's, in a manner inconsistent with its labeling.

Observe the following precautions when mixing:

- Make sure all spray and application equipment are clean prior to mixing Arroz 80; clean equipment well after completing application of Arroz 80 (see **PREPARATION AND CLEANUP OF APPLICATION EQUIPMENT**, below)
- Do not allow Arroz 80 packets to become wet prior to mixing, and do not handle packets with wet gloves
- If any Arroz 80 packets are unused, outer container must be closed and tightly resealed to protect the packets and preserve the integrity of the water soluble packaging
- Make sure that water soluble packets have completely dissolved prior to adding any additional ingredients (it should take the packets about 5 minutes to wholly dissolve)
- Cold water, insufficient agitation or water with high rates of sulfur or boron could unfavorably affect dispersal of Arroz 80, resulting in potential clogging of nozzle or spray screen

Arroz 80 can be kept in the mix or spray tank for three days following mixing, without a reduction in efficacy. If spray solution is held for a period of time, be sure to mix/agitate fully prior to use.

PREPARATION AND CLEANUP OF APPLICATION EQUIPMENT

PRECAUTION: DO NOT USE chlorine bleach for cleaning or mix chlorine bleach with ammonia. Make certain that all traces of any fertilizer containing ammonia or ammonium are completely removed before adding any chlorine (including chlorine bleach) to the mix tank.

Adverse crop reaction may result if residues of previously applied products are left in application equipment, or if residues of Arroz 80 are left in spray equipment following application. Clean spray equipment prior to using Arroz 80, and clean immediately after treatment with Arroz 80, and before applications with other products.

Before using Arroz 80, completely drain, rinse and clean all spray and mixing equipment, following procedures instructed for the previously used product. If previously sprayed product is not completely removed, Arroz 80 residues could collect in the spray equipment resulting in clogged equipment or greater difficulty in cleaning after use of Arroz 80

After spraying Arroz 80, use the following procedure to clean equipment:

1. Remove any visible residue
2. Drain the spray application equipment, including tank, hoses, spray boom and nozzles.
3. Fill tank 50% full of water, spraying the interior sides of the tank while filling
4. Use a tank cleaner that DOES NOT contain chlorine, and fill the remainder of the tank with clean water. Follow tank cleaner instructions regarding agitation/recirculation of the cleaner throughout the tank, boom and hoses; completely flush boom and hoses prior to draining the tank

5. Rinse with clean water to remove tank cleaner from tanks, boom, hoses, nozzles and strainers (follow any directions provided with tank cleaner)
6. Fill tank 50% full of water, and add 3% active household ammonia (1 gallon per every 100 gallons tank size). Finish filling the tank with clean water, and recirculate the ammonia solution for 15 minutes; completely flush tank, boom, hoses, nozzles and strainers prior to draining the tank
7. Remove strainers, screens and nozzles, and clean independently in a solution of 3% active household ammonia and water, then replace all strainers, nozzles and screens
8. Repeat step 6 (ammonia cleaning step)
9. Completely rinse tank and equipment with clean water, and flush clean water through hoses, boom and nozzles so that all ammonia is removed

Dispose rinse solution at an approved waste disposal location or on-site.

**USE DIRECTIONS: DRY-SEEDED OR WATER-SEEDED RICE – U.S. RICE GROWING REGIONS
(Except California)**

Arroz 80 can be applied in the following use patterns, either by itself or as a tank mix partner (see **TANK MIXES** section, below):

- Single Arroz 80 application (solo or tank mix)
- Early postemergence Arroz 80 application (tank mixed with preemergence herbicide) followed by pre- or post- flood Arroz 80 application (solo or tank mix)
- Mid postemergence Arroz 80 application (solo or tank mix) followed by pre-or post-flood Arroz 80 application

Single Application -- See **WEEDS AND USE RATES** chart for rates and timings and weeds controlled
Early Postemergence -- When rice has reached the 2-leaf growth stage (when 2nd leaf is fully expanded), make first application of Arroz 80 at 0.2 oz. / A tank mixed with a rice preemergence herbicide containing the active ingredients thiobencarb (including Bolero® 8 EC), clomazone (including Command® 3ME), quinclorac (including Facet®) or pendimethalin (including Prowl® 3.3 EC) -- see **TANK MIX** section, below, and check tank mix partner label for specified use rate. Make second application of Arroz 80 at 0.53 to 0.67 oz./A just before permanent flood, or early post-flood (see **WEEDS AND USE RATES** chart).

Mid Postemergence -- When barnyardgrass reaches the 3- to 5-leaf growth stage, make first application of Arroz 80 at 0.5 oz. / A. Make second application of Arroz 80 at 0.5 oz./a just before permanent flood, or early post-flood (see **WEEDS AND USE RATES** chart).

WEEDS AND USE RATES FOR USE IN RICE GROWING REGIONS (EXCEPT CALIFORNIA)			
Weed	Weed Size	Control or Suppression	Use Rate (oz. / A)
Alligatorweed (<i>Alternanthera philoxeroides</i>)	Up to 10 inch runners	S	0.53-0.57
Annual Rice Flatsedge (<i>Cyperus iria</i>)	1 -3 tillers	C	0.57-0.67
Barnyardgrass / Junglerice ¹ (<i>Echinochloa crus-galli</i> / <i>Echinochloa colona</i>)	2-leaf up to 5 leaf	C	0.4
	5 leaf through 1 tiller	C	0.53
	Up to 3 tillers	C	0.57
Barnyardgrass / Junglerice (<i>Echinochloa crus-galli</i> / <i>Echinochloa colona</i>) – Late Application ²		S	0.57-0.67
Barnyardgrass, perennial (<i>Echinochloa polystachya</i>)	Up to 2 tillers	S	0.53-0.57
Baronet grass (bayonetgrass) – (<i>Echinochloa pungens</i>) – Post Flood Only	1 to 3 tillers	C	0.57-0.67
Dayflower (<i>Commelina communis</i>)	1 leaf up to 4 leaf	C	0.4-0.57

Ducksalad (<i>Heteranthera</i> spp.)	1 leaf up to 4 leaf	C	0.4-0.57
Eclipta (<i>Eclipta</i> spp.)	1 leaf up to 4 leaf	S	0.4-0.57
Gooseseed (<i>Sphenoclea zeylanica</i>)	1 leaf up to 4 leaf	C	0.4-0.57
Hemp Sesbania (<i>Sesbania exaltata</i>)	3 to 18 inches	C	0.4-0.57
Johnsongrass (<i>Sorghum helepense</i>)	3 to 24 inches	C	0.4-0.57
Jointvetch, Indial (<i>Aeschynomene indica</i>)	3 to 18 inches	C	0.4-0.57
Jointvetch, Northern (<i>Aeschynomene virginica</i>)	3 to 18 inches	C	0.4-0.57
Knotgrass (<i>Paspalum ditichum</i>) – Post Flood Only ³	Up to Heading	S	0.53-0.57
Morningglory, entireleaf (<i>Ipomoea hederacea</i>)	1 to 4 inches	S	0.4-0.57
Morningglory, pitted (<i>Ipomoea lacunose</i>)	1 to 4 inches	S	0.4-0.57
Pigweeds (<i>Amaranthus</i> spp.)	1 to 12 inches	S	0.4-0.57
Redstem (<i>Ammannia</i> spp.)	1 to 4 inches	S	0.4-0.57
Smartweed, Pennsylvania (<i>Polygonum pennsylvanicum</i>)	1 to 4 inches	C	0.4-0.57
	4 to 24 inches	S	0.4-0.57
Texas / Mexicanweed (<i>Caperonia</i> spp.)	1 leaf up to 4 leaf	S	0.4-0.57
Water Hyssop (<i>Bacopa rotundifolia</i>)	1 leaf up to 4leaf	C	0.4-0.57

¹ Includes propanil and or Facet (quinclorac) resistant barnyardgrass

² If barnyardgrass reaches the 4-tiller up to booting growth stages, it has begun to adversely affect rice yields. Suppression or control at this time will be beneficial by reducing production of barnyardgrass seed, and by making the most of remaining rice yield

³ For best results in suppressing knotgrass, apply before knotgrass heading, after rice is in permanent flood, when a minimum of 70% of the knotgrass is above the water level.

- When making an early postemergence split application, make application to rice that has reached the 2-leaf growth stage (2nd leaf fully expanded) or after panicle initiation growth stage (green ring appears, just before joint movement) at the lower specified use rate
- For all other applications, do not apply to rice until it has reached the 3-leaf growth stage (3rd leaf fully expanded) – irrespective of seeding method - with a root system totally underneath soil surface. Application can be made up to the point of panicle initiation (green ring appears, just before joint movement).
- After application of Arroz 80, rice plants may exhibit temporary chlorosis, stunting or other injury. This injury is not permanent, and rice plants will recover. Top dressing with fertilizer can hasten recovery.
- If rice is not fully pegged (root system totally underneath soil surface), application of Arroz 80 could result in considerable injury, despite growth stage.
- **Pre-Flood Application** - When applying Arroz 80 pre-flood, optimum results are obtained when soil is wet to the surface and weeds are actively growing. Allow herbicide at least one day for uptake after application before establishing the permanent flood. If permanent flood is delayed (to allow rice to become tolerant to flood), flush as required to support rice growth and weed growth (which, in turn, supports herbicide uptake). Herbicidal efficacy can be compromised if soil becomes dry after application of Arroz 80. For best results, establish permanent flood 2 to 7 days after application of Arroz 80. Weed reinfestation and/or reinvigorated growth of existing weeds can result if permanent flood is held off too long.
- **Post-Flood Application** - When applying Arroz 80 post-flood, optimum results are obtained when flood water is adjusted so that a minimum of 70% of the weed plant is above the water level. 2 to 3 days after treatment, water level can be raised to normal flood level.
- For best results make application of Arroz 80 when nighttime temperatures have been at 60° F or higher for at least 3 consecutive nights before application. Lower nighttime temperatures can result in reduced herbicidal efficacy.
- Rice under stress due to environmental conditions (drought, temperature, etc.) or other conditions (nutrient deficiencies or injury due to herbicide or fertilizer applications) which reduce the plant's metabolism and

development can exhibit sensitivity to Arroz 80. Likewise, weeds under similar stress will not be as susceptible to Arroz 80 treatment. Do not apply to stressed rice or weeds.

- Medium grain rice varieties, and pubescent (hairy) leaf rice varieties may exhibit more sensitivity to Arroz 80 than long grain or glabrous (smooth) leaf rice varieties. Rice varieties with low seedling vigor (including M-206 or Japanese cultivars) may exhibit sensitivity to Arroz 80, particularly if they are under environmental or other stress. Do not apply Arroz 80 to Bengal rice variety.
- Arroz 80 can be applied to hybrid varieties of rice, including Clearfield® rice
- When a use rate range is given for a particular weed species, use the upper end of the specified rate range if weed infestation is elevated or if weeds are approaching upper end of specified weed size. If infestation is severe, a second application of Arroz 80 or another herbicide may be required for control.
- Growers can make additional applications of Arroz 80, as long as the maximum seasonal application rate of 1.06 oz. product per acre and application interval of 3 weeks are observed.

ADDITIVES

Surfactants - Apply Arroz 80 with a surfactant, unless specific label section or supplemental label indicates otherwise. See 'Arroz 80 Approved Surfactants' bulletin for a list of permitted surfactants and use rates. Use of any surfactant other than those indicated in the approved surfactants bulletin is done at the sole discretion and risk of the user.

Urea-ammonium Nitrate (UAN) – If chosen surfactant does not already contain UAN, addition of 2% volume/volume of 28% to 32% UAN, in addition to an approved surfactant can heighten the efficacy of Arroz 80.

TANK MIXES

For broader weed spectrum control, Arroz 80 may be used in combination with other herbicides. It is the pesticide user's responsibility to ensure that all products are registered for intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank mix partners for Arroz 80 include products containing the following herbicide active ingredients:

2,4-D	Pendimethalin (including Prowl 3.3 EC)
Bensulfuron methyl (including Londax®)	Quinclorac (including Facet)
Clomazone (including Command 3ME)	Sodium acifluorfen (including Blazer® or Ultra Blazer)
Diflufenzuron (including Cavalier 2L or Dimilin®)	Thiobencarb (including Bolero 8 EC)
Fenoxaprop-p-ethyl (including Ricestar® HT)	Triclopyr (including Grandstand® R)
Halosulfuron-methyl (including Permit®)	

Arroz 80 can also be tank mixed with Clearpath® or Newpath® herbicides in Clearfield rice only.

Arroz 80 can also be tank mixed with other pesticides, including those containing the insecticide active ingredients lambda cyhalothrin (including Karate® or Karate Z) or zeta-cypermethrin (including Fury®); or fungicide active ingredient azoxystrobin (including Quadris®).

Take care when tank mixing Arroz 80 with products containing the herbicide active ingredient carfentrazone-ethyl (including Aim®). Carfentrazone ethyl can result in antagonism to bispyribac sodium activity, and may result in the need for an additional application of Arroz 80 or other herbicide. If applying an Arroz 80 – carfentrazone ethyl tank mix, go up to the next Arroz 80 use rate for the particular weed size, and limit use rate of Aim to no more than 1 fl. oz. product per acre (please refer to Aim label for current labeled use rate on rice – if it is lower than 1 fl. oz. per acre, DO NOT EXCEED the labeled rate).

When tank mixing Arroz 80 with quinclorac, use adjuvants/surfactants instructed for Arroz 80, and do not include a crop oil concentrate.

Not all rice varieties have been tested with all possible tank mix combinations. If you are not familiar with an Arroz 80 tank mix with any of the listed products, or a tank mix with a pesticide product that is not listed in this section, it is your responsibility to test the combination for crop safety on a small portion of your rice crop to ensure that a phytotoxic or other adverse response will not occur. In addition, test the physical compatibility of Arroz 80 with tank mix partners before use. In a lidded glass jar (~1 quart size), add all mix partners, in their relative proportions. Invert, shake or mix the jar thoroughly. Observe mixture for approximately 30 minutes (though signs of incompatibility will often be seen within 5 minutes).

Tank Mix Restrictions:

- To avoid injury or antagonism, do not tank mix Arroz 80 with pesticide products containing the active ingredients malathion, methyl parathion or propanil
- Do not apply Arroz 80 within 7 days of treatment with malathion or methyl parathion

USE RESTRICTIONS

- Do not irrigate other crops with water that has been drained directly from fields treated with Arroz 80
- Field ends cannot be double-sprayed
- Apply a maximum 1.06 oz. Arroz 80 per acre per year. Do not make application to second crop (stubble/ratoon crop) rice.
- Observe a 3-week application interval for subsequent applications of Arroz 80
- Do not treat stressed rice or weeds with Arroz 80
- If fields have been land leveled and have extreme cut and heavy fill areas, Arroz 80 cannot be applied to the first rice crop in these fields (this restriction does not pertain to maintenance leveling).
- If commercial crayfish farming is practiced in rice paddies, Arroz 80 cannot be applied.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool dry place. Keep pesticide in original container. Keep container closed when not in use. Do not put concentrate or dilute into food or drink containers. Not for use or storage in or around the home. For help with any spill, leak, fire, or exposure involving this material, call day or night 800-424-9300.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

For outer bag containing water soluble packets

Nonrefillable outer bag. Do not reuse or refill the outer bag. Offer for recycling, if available or dispose of outer bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2020 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMERICAN VANGUARD, AMVAC, ARROZ, ARROZ 80, and their respective logos, are trademarks owned by AMVAC Chemical Corporation.

CHEMTREC is a service mark of the American Chemistry Council, Inc. Viton is a trademark of the Chemours Company FC, LLC. Bolero is a trademark of Kumiai Chemical Industry Co., Ltd. Aim, Command, and Fury are trademarks of FMC Corporation. Blazer and Ultra Blazer are trademarks of United Phosphorus Inc. Clearfield, Clearpath, Facet, Newpath, and Prowl are trademarks of a BASF company. Dimilin is a trademark of Arysta LifeScience, Inc. Grandstand is a trademark of Dow AgroSciences, LLC. Karate and Quadris are trademarks of Syngenta Ltd. Londax is a trademark of United Phosphorus Ltd. Permit is a trademark of Nissan Chemical Corp. Ricestar is a trademark of Bayer Crop Science Ag.

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.