11



EPA Reg. No. 91234-74

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FIFRA SECTION 2(EE) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALL STATES EXCEPT ALABAMA, ARKANSAS, NEW YORK, SOUTH CAROLINA AND TEXAS

For Control of Fairy Ring on Turf

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **Atticus Artavia™ 2 SC.**

TURF

Golf course turf (not for use in California) Commercial turf farms (not for use in California)

Atticus Artavia™ 2 SC is specified for control on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Application Instructions*
Fairy Ring (Lycoperdon spp., Arachnion spp., Bovista spp., Vascellum spp., and Agrocybe pediades)	0.38 – 0.77	14 - 28	Apply preventatively or as soon as possible after Fairy ring symptoms develop. Apply only in 4 gals. of water per 1000 sq. ft. (174 gals. / Ac.). Add the specified rate of a wetting agent to the final spray. Fairy ring symptoms may take 2 to 3 weeks to disappear following curative applications and reapplication may be required in some cases. Severely damaged or thin Turf may require reseeding.

*Do not apply more than 4 sequential applications of Atticus Artavia ™ 2 SC

