



**24(c) Special Local Need Label
For Use on Horseradish**

EPA SLN No.: IL-140001
EPA Reg. No. 279-3442

**FOR DISTRIBUTION AND USE
ONLY IN ILLINOIS**

This label for Authority Elite Herbicide is valid until April 28, 2019, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

DIRECTIONS FOR USE

CROP/SITE:	Weeds Controlled		RATE:
Horseradish	barnyardgrass fall panicum foxtail, giant foxtail, green foxtail, yellow morningglory, entireleaf morningglory, ivyleaf morningglory, pitted morningglory, smallflower	nightshade, black nightshade, eastern black palmer amaranth Pennsylvania smartweed pigweed, red root pigweed, smooth waterhemp, common waterhemp, tall	19 to 25 fluid ounces per acre

Apply in at least 10 gallons per acre finished spray solution by ground.

Apply a single application of Authority® Elite at a broadcast rate of 19-25 fluid ounces per acre to the soil surface after planting but before weed or crop emergence. Use lower rates on soils relatively coarse-textured and higher rates on fine textured soils.

Following the application of Authority Elite herbicide to soil, germinating seeds and seedlings take up Authority Elite from the soil solution. The amount of Authority Elite herbicide in soil solution available for weed uptake is determined primarily by soil type, soil organic matter and soil pH. Similar to other herbicides, Authority Elite herbicide adsorbs to the clay and organic matter (OM) fractions of soils; effectively limiting the amount of active ingredient immediately available to control weeds. Adequate moisture is required for herbicide activation (1/2" to 1" of rainfall or irrigation). If an activating rainfall (1/2" to 1") is not received Authority Elite herbicide will provide a reduced level of control of susceptible germinating weeds.

RESTRICTIONS:

Do not exceed 25 fluid ounces per acre per cropping season.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply directly on the crop after the crop emerges or if the seedling sprouts are close to the soil surface.

Harvest horseradish at normal timing.

Label code 043014
 Copyright 2014. All rights reserved.
 Authority is a trademark of FMC Corporation.

FMC
 FMC Corporation
 Agricultural Products Group
 1735 Market Street
 Philadelphia, PA 19103