



FIFRA §2(ee) RECOMMENDATION
For distribution and use only within
the states of Connecticut, Delaware,
Maryland, Massachusetts,
New Hampshire, New Jersey,
Pennsylvania, Rhode Island,
Vermont and Virginia.



Controlled Release Insecticide

EPA Registration number 74530-53-5481

This recommendation is made as permitted under FIFRA Section 2(ee) and therefore has not been submitted to or approved by the Environmental Protection Agency (EPA).

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label

DIRECTIONS FOR USE

Refer to the Avesta CS label for complete Directions for Use paying attention to use site descriptions.

OUTDOOR SURFACES USE

For control of Spotted Lanternfly nymphs and adults

Apply with either hand or power application equipment as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages, refuse dumps, and other similar areas where these insect pests are active.

Application Rate: Fl. oz. (mL) of AVESTA CS	Gals. of Water¹	Area of Coverage (sq. ft.)
0.2 fl. oz. (6mL)	1-5 gals.	800-1600 sq. ft.
0.4 fl. oz. (12 mL)	1-5 gals.	800-1600 sq. ft.
0.8 fl. oz. (24 mL)	1-5 gals.	800-1600 sq. ft.

¹Application volume may be greater than 5 gals./800-1600 sq. ft. if required under heavy vegetative or landscaping materials in order to obtain desired coverage.

INSECT CONTROL ON ORNAMENTAL PLANTS

AVESTA CS is for use to control the pest in such areas as residential landscaped areas and landscaped areas around institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields (including trees, shrubs, flowers, evergreens, foliage plants and groundcovers). Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at recommended rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. Apply at 7-day intervals if retreatment is necessary.

Limit more frequent treatment to spot treatments. The higher application rate should be used when maximum residual control is desired.

Pest	Application Rate Avesta CS	
	Spot Treatment: Fluid Ounces per 1 Gallon	Standard Treatment: Fluid ounces per 100 gallons
Spotted Lanternfly nymphs and adults	0.2 to 0.5	1.5 to 5

RESTRICTIONS

- Let treated surfaces dry before allowing humans and pets to contact surfaces.
- DO NOT use this product with oil.
- DO NOT apply to institutions (including libraries, sports facilities, etc.,) in the immediate area when occupants are present.
- DO NOT apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- DO NOT allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are located in/around structures being treated.
- DO NOT apply this product through any type of irrigation system.
- DO NOT apply this product by aerial application.
- DO NOT apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.
- DO NOT make outdoor broadcast applications to ornamentals when wind speed is 15 mph or greater.
- DO NOT apply more than 0.36 lbs. of the a.i. (52.4 fl. oz. of concentrate) per acre per year.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer before using this product.

AMVAC, the AMVAC logo, and Avesta are trademarks owned by AMVAC Chemical Corporation.

Registrant: AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200,
Newport Beach, CA 92660
1-888-462-6822