



Axcel™

0-0-1

Guaranteed Analysis:

Soluble Potash (K ₂ O).....	1%
Calcium (Ca).....	1%
Manganese (Mn).....	0.4%
0.4% Water Soluble Manganese (Mn)	
Molybdenum (Mo).....	0.017%

Derived from: Rock Phosphate,
manganese sulfate, sodium molybdate, kelp

Net Contents:

<input type="checkbox"/> 1 gallon (3.78L) 9.26 lbs (4.2 kg)	<input type="checkbox"/> 55 gallon (208.2L) 509.3bs (231 kg)
<input type="checkbox"/> 5 gallon (18.92L) 46.3 lbs (21 kg)	
<input type="checkbox"/> 30 gallon (113.56L) 277.8 lbs (126 kg)	<input type="checkbox"/> 275 gallon (1041L) 2546.5 lbs (1155 kg)

F2315

Density: 9.26 lbs. per gallon at 68°F

Information regarding the contents and levels of
metals in this product is available on the internet at
<http://www.aapfco.org/metals.html>

Disclaimer: AEA products are sold "as is" and all warranties, express or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules and other international bio-organic regulations for certified organic crop production. Verify with your certifier before using

PURPOSE: Axcel™ is a nutritionally-based reproductive support package.

APPLICATION RATES AND TIMING: This product is intended for foliar and fertigation applications and should only be used in accordance with the manufacturer's instructions. Check with your crop consultant for specific application rates. Do not harvest within 14 days of application.

Single Flowering Crops: Use a single foliar application of up to 4 gal. per acre 1-2 weeks before flowering or tuber-set commences. For fertigation, apply up to 4 gal. per acre in 2 applications within the two weeks before flowering.

Continuously Flowering Crops: Use an initial foliar application of up to 2 gal. per acre 1-2 weeks before flowering commences and up to 4 quarts per acre every two weeks until flowering is complete. Use the same rates for drip fertigation.

CAUTION: The application of fertilizers containing molybdenum may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

Handling: Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When tank mixing: A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however if they are inadequately diluted or agitated before and during application there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.



Advancing Eco Agriculture, LLC

4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603