



GeaPower is the exclusive technology platform developed by Valagro


**KEEP OUT OF REACH OF CHILDREN  
CLEANING OF SPRAYER**

Flush with clean water all parts of the sprayer including the pump, hoses and nozzles several times prior to and after the use of this product.

**CONDITIONS OF SALE**

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions and use. No one other than an officer of the seller is authorized to make any warranty, guarantee or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller's control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price. It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions, abnormal conditions, presence of other materials, the manner of application, or other factors, all of which are beyond the control of Valagro USA or the seller. All such risks shall be assumed by buyer.

**AXILO RMX**



**WARNING**  
H319 Causes serious eye irritation.  
Wash hands thoroughly after handling.  
Wear protective safety goggles and face shield  
IF IN EYES: Rinse cautiously with water for several minutes.  
Remove contact lenses, if present and easy to do.  
Continue rinsing. If eye irritation persists: Get medical advice/attention.

**FIRST AID:** As a general rule, in case of doubt or if symptoms persist, always call a doctor.  
Inhalation: Remove to fresh air.  
Get medical attention for any breathing difficulty.  
Eye Contact: Rinse cautiously with water for several minutes holding the eyelids open. Remove contact lenses, if present and easy to do. Continue rinsing. If irritation occurs and if symptoms persist, call a physician.  
Ingestion: Rinse mouth with water. Induce vomiting only if indicated by the doctor. Never give anything by mouth to an unconscious person. Immediately call a doctor.  
Skin Contact: Remove all contaminated clothing. Rinse with plenty of soap and water.

Produced and packed by VALAGRO S.p.A.  
Zona Ind.le - 66041 Atessa (CH) - ITALY- Tel. (+39) 0872 / 8811  
Distributed and guaranteed by Valagro USA Inc.  
2020 Ponce de Leon Blvd, Ste. 1203 - Coral Gables, FL 33134  
Phone +1-786-230-1020 / +1-786-230-1019 / Fax +1-786-475-9032  
Code: see code on the package  
Lot number: see print on package

**NET WEIGHT 20 Lb (9.07 kg)**

**DIRECTIONS FOR MIXING**

Add half a pound of Axilo to one gallon of water to make a slurry which can be then added to starter solutions or tank mixes. Add buffer to solutions over pH 6.5. Dissolve product into a convenient amount of water, consistent with the type of sprayer used. Choose a water volume necessary to achieve uniform coverage with little or no run-off. Spray nozzles should produce a fine mist. Spraying should be carried out on a calm day but not during strong sunshine. Early morning or late afternoon applications are best. If rainfall occurs within several hours after application, the crop may require respraying 3 to 4 days later.

**TANK MIXES**

If the product is to be applied alone, a non-ionic wetting agent should be added to the spray tank in accordance with the manufacturer's direction. If applied in conjunction with chemicals which contain their own wetting agent, then the addition of a wetting agent may not be necessary. If using in conjunction with a hormone type selective weed killer, consult the manufacturer of the effect of wetting agents on their herbicide (many wetting agents tend to destroy the selectivity of hormone type weed killers).



www.valagro.com

AXILO RMX | Valagro®

002574

# AXILO RMX

**SOLUBLE MICROGRANULES  
GUARANTEED ANALYSIS**

Copper (Cu).....	2.6%
2.6% chelated Copper (Cu)	
Iron (Fe) .....	5.6%
5.6% chelated Iron (Fe)	
Manganese (Mn) .....	2.6%
2.6% chelated Manganese (Mn)	
Zinc (Zn) .....	2.6%
2.6% chelated Zinc (Zn)	
Derived from Copper EDTA, Iron EDTA, Manganese EDTA and Zinc EDTA	

**GENERAL INFORMATION**

Axilo RMX is compatible with a wide range of fertilizers and pesticides, however, a compatibility jar test should be performed before mixing, especially with phosphorous fertilizers.

**20lb NET WEIGHT (9,07 Kg)**

Distributed and guaranteed by Valagro USA Inc.  
2020 Ponce de Leon Blvd, Ste. 1203 - Coral Gables, FL 33134  
Phone +1-786-230-1020 / +1-786-230-1019 / Fax +1-786-475-9032

FOLIAR APPLICATION



ROOT APPLICATION



Where science serves nature

MADE IN ITALY

AXILO RMX | Valagro®

**| AXILO RMX**

**DIRECTIONS FOR USE**

(Use low rates for maintenance applications, middle rates for medium deficiencies and high rates for extreme deficiencies)

**SOIL APPLICATION**

Can be used with starter fertilizers but mixture compatibility is unpredictable and is at the user's risk. Jar testing is always recommended along with use of sufficient water to slurry the Axilo. Surface application by sprayer and cultivated into the soil immediately after spraying. Mixed into a slurry and impregnated onto granular fertilizer. See "Directions for Mixing" section.

**FOLIAR APPLICATION**

Can form a stable mixture with glyphosate, other pesticides and foliar fertilizers. See sections on "Directions for Mixing" and "Tank Mixes"  
DO NOT apply this product when the crop is under water stress. Application after irrigation or rainfall is best.  
DO NOT apply to plants bearing mature fruit until it has been determined that this product will not damage or mark the fruit in any way.

**VEGETABLE CROPS:** Including but not limited to: onion, carrot, pepper, eggplant, zucchini, melon, water melon, cucumber, tomatoes, cole crops, leafy vegetables, etc.; 0.25 - 2.0 lb per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**TREE & VINE FRUIT CROPS:** Including but not limited to: apples, cherries, citrus, pear, stone fruit, kiwi, grapes (raisin, table, wine) etc.; 2-8 oz per tree or, 0.25 - 2.0 lb per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**SHRUB, BERRY AND BUSH CROPS:** Including but not limited to: blueberries, cranberries, raspberries, strawberries etc.; 2-4 lb per 100 bushes or, 0.25 - 2.0 lbs per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**NUT CROPS:** Including but not limited to: almond, pecan, pistachio, walnut, etc.; 2-8 oz per tree or, 0.25 - 2.0 lb per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**FIELD ROW CROPS:** Including but not limited to: barley, beans (edible, field, snap, etc), corn (field, silage, seed, sweet, etc), cotton, peanuts, potatoes, rice, soybeans, sorghum (grain, sweet, etc), sugar cane, sugar beets, wheat etc.; 0.25 - 2.0 lb per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**FORAGE/GRASS CROPS:** Including but not limited to: alfalfa, timothy, fescue, switchgrass etc.; 0.25 - 2.0 lb per broadcast acre; Apply maintenance (lower) during vegetative growth and then, 14-21 days until deficiency symptoms disappear.

**GOLF COURSE GREENS:** ½ lb/acre (43,650 sq ft) in a minimum, 150-200 gallons of water; Apply in early spring when greens are actively growing. Four or more applications can be made at monthly intervals to maintain color and vigor. Applications should always be made late in the evening or early in the morning to reduce potential for phytotoxicity.

**TURF:** Including but not limited to: sod, lawns, golf course fairways and tees, playing fields, recreational areas etc.; 0.25 - 2.0 lb per acre in a minimum, 40 gallons of water; 3 or more applications per year to maintain vigorous turf.

**ORNAMENTALS:** 0.25 - 2.0 lb per broadcast acre; Apply as a drench between rows and cultivate into the soil early in the season. Irrigate after application.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Do not apply near water, storm drains, or drainage ditches.  
Do not apply if heavy rains are expected.  
Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.



8 031702 005668

4242267F07P159 02