

BIOMIN[®] Mn & Zn

2-0-0

Chelated Mineral for Soil and Foliar Application



Product Information

BIOMIN Mn & Zn contains chelated manganese and zinc for soil and/or foliar application. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct manganese and zinc deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	2.0%
2.0% Water Soluble Nitrogen	
Manganese (Mn).....	5.0%
5.0% Chelated Manganese	
Zinc (Zn).....	6.0%
6.0% Chelated Zinc	

Derived from manganese sulfate, zinc sulfate, citric acid (chelating agent), and hydrolyzed vegetable protein.

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

**PREVENTS AND CORRECTS
MANGANESE AND ZINC DEFICIENCIES**

NET CONTENTS: 2.5 gal. (9.45 l)

Weight per gallon: 10.8 lb. (4.9 kg) @ 68 °F

U.S. Pat.# 5,504,055

Lot #:

**Made in
USA**

Directions and General Recommendations

SOIL APPLICATION

BIOMIN Mn & Zn can be applied before or at planting as banding or broadcast. The application rate is 1 qt. to 1 gal./ac. (2.5-10 l/ha) depending on soil conditions and application methods. For maintenance, 1 qt./ac. (2.5 l/ha) is recommended. Dilute **BIOMIN Mn & Zn** with enough water to cover the entire acre.

FOLIAR APPLICATION

FIELD CROPS: Apply 1-4 pt./ac. (1.5-5 l/ha) of **BIOMIN Mn & Zn** when crops are about 4-6 in. (10-20 cm) high or when deficiency symptoms appear. Repeat the application in 2-4 weeks if necessary.

FRUIT TREES AND GRAPES: Apply 1-4 pt./ac. (1.5-5 l/ha) of **BIOMIN Mn & Zn** starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

NUT TREES: Apply 1-4 pt./ac. (1.5-5 l/ha) of **BIOMIN Mn & Zn** starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

VEGETABLE CROPS: Apply 1-4 pt./ac. (1.5-5 l/ha) of **BIOMIN Mn & Zn** at about 2-4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1-3 weeks if necessary.

For best results, apply **BIOMIN Mn & Zn** when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, dilute it with at least 100 times of water. **BIOMIN Mn & Zn** will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN Mn & Zn can be applied to strawberry, lettuce, tomato, celery, grapes, stone fruits, tree fruits, vegetables, commercial turf, and ornamental gardens.

Precautionary statement: do not swallow. Wash hands thoroughly after handling. **First Aid:** if swallowed: Seek medical attention if feeling unwell. Rinse mouth. **Storage & Disposal:** store in a safe manner. Dispose product/container in accordance with federal, state, and local regulations.

JHBiotech, Inc.

0345-CA-B-1



MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538 , Ventura, CA 93006 U.S.A. 805-650-8933 jhbiotech.com

Copyright © 2016 All rights reserved