

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

BAYTHROID® XL

For the control of Boll weevil in Sugarcane and Corn

EPA Reg. No. 264-840

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF TEXAS

This recommendation is valid until December 31, 2024

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

SUGARCANE		
PESTS CONTROLLED	Rate fluid oz/Acre	Rate lb AI/Acre
Boll Weevil	2.8	0.022
Foliar Application Restrictions Pre-Harvest Interval (PHI): 15 days . Maximum BAYTHROID XL allowed per 7-day interval: 2.8 fluid oz/A (0.022 lb AI/Acre) . Maximum BAYTHROID XL allowed per crop season: 16.8 fluid oz/A (0.132 lb AI/Acre) . For ground application, apply in a minimum of 10 GPA. Minimum ULV application volume (once refined cotton seed/vegetable oil): 1.0 qt/A – aerial application. Do not apply if soil is saturated with water. Do not apply under conditions that favor runoff. Do not apply in the rain.		

CORN – Foliar Applications**Field Corn, Popcorn, Seed Corn, Teosinte – (see Sweet Corn application information in Vegetable Crops Section)**

PESTS CONTROLLED	Rate fluid oz/Acre	Rate lb AI/Acre
Boll Weevil	2.8	0.022

Foliar Application RestrictionsPre-Harvest Interval (PHI): Grain or fodder – **21 days**; Green forage – **0 day**.Maximum BAYTHROID XL allowed per 7-day interval: **2.8 fluid oz/A (0.022 lb AI/Acre)**.Maximum BAYTHROID XL allowed per crop season: **11.2 fluid oz/A (0.088 lb AI/Acre)**.Maximum number of applications per season: **4**. Three applications may be applied up to early dent stage. One application may be made between early dent and 21 days before harvest.

Minimum ULV application volume (once refined cotton seed/vegetable oil): 1.0 qt/A – aerial application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).*As with any crop-protection product, always read and follow label directions.**For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*