

# RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

## BAYTHROID® XL

To add the control of the pest Wireworm to existing 2(ee) for the control of Cabbage Maggot, Garden Symphylan and Seed Corn Maggot in Brassica and Leafy Vegetables.

EPA Reg. No. 264-840

### *FIFRA 2(ee) Recommendation*

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CALIFORNIA

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### BRASSICA (COLE) LEAFY VEGETABLES

Includes all members of Crop Group 5:

Broccoli, Broccoli raab (rapini), Chinese (gai lon) broccoli, Brussels sprouts, Cabbage, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, and Turnip greens.

PESTS CONTROLLED	Rate fluid oz/Acre	Rate lb AI/Acre	Application Method
Cabbage Maggot Garden Symphylan Seed Corn Maggot Wireworm	3.2	0.025	For reduction of damage caused by the listed pests, apply specified dosage per acre as a soil-incorporated treatment before or at planting or transplanting.

#### Notes and Restrictions

Maximum number of pre-plant or at-plant/transplant applications: 1

Maximum BAYTHROID XL allowed per 7-day interval: **3.2 fluid oz/A (0.025 lb AI/Acre)**.

Maximum BAYTHROID XL allowed per crop season: **12.8 fluid oz/A (0.1 lb AI/Acre)**.

Refer to product label for application instructions and guidelines.

**LEAFY VEGETABLES**

Includes all members of Crop Group 4:

**Amaranth (Chinese spinach), Arugula (rouquette), Cardoon, Celery, Chinese celery, Celtuce, Chervil, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Endive (escarole), Florence fennel, Lettuce (head and leaf), New Zealand spinach, Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, Swiss chard, Vine spinach**

<b>PESTS CONTROLLED</b>	<b>Rate fluid oz/Acre</b>	<b>Rate lb AI/Acre</b>	<b>Application Method</b>
Cabbage Maggot Garden Symphylan Seed Corn Maggot Wireworm	3.2	0.025	For reduction of damage caused by the listed pests, apply specified dosage per acre as a soil-incorporated treatment before or at planting or transplanting.

**Notes and Restrictions:**

Maximum number of pre-plant or at-plant/transplant applications: 1

Maximum BAYTHROID XL allowed per 7-day interval: **3.2 fluid oz/A (0.025 lb AI/Acre)**.Maximum BAYTHROID XL allowed per crop season: **12.8 fluid oz/A (0.1 lb AI/Acre)**.

Refer to product label for application instructions and guidelines.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).*As with any crop-protection product, always read and follow label directions.**For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*