


**RESTRICTED USE PESTICIDE**  
**Due to Toxicity to Fish and Aquatic Organisms**  
**For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.**

 <b>Bayer CropScience LP</b> <b>P.O. Box 12014</b> <b>2 T.W. Alexander Drive</b> <b>Research Triangle Park, North Carolina</b> <b>27709</b> <b>1-866-99BAYER (1-866-992-2937)</b>	<table border="1"> <tr> <td><b>GROUP</b></td> <td style="background-color: #cccccc;"><b>3</b></td> <td><b>INSECTICIDE</b></td> </tr> </table>	<b>GROUP</b>	<b>3</b>	<b>INSECTICIDE</b>
	<b>GROUP</b>	<b>3</b>	<b>INSECTICIDE</b>	
<b>Baythroid® XL</b> <b>EPA Reg. No. 264-840</b>  <b>For use on Grape, Potato and other</b> <b>Tuberous and Corm Vegetables to control</b> <b>Brown Marmorated Stink Bug</b>				

## ***PRODUCT BULLETIN***

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF**  
**DELAWARE, MARYLAND, NEW JERSEY, PENNSYLVANIA, VIRGINIA, and WEST VIRGINIA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product inconsistent with its labeling.

<b>GRAPE</b>		
Includes: Table grape, Raisin, Wine and Muscadine grape		
<b>PEST CONTROLLED</b>	<b>Rate fluid ounces/Acre</b>	<b>Rate lb AI/Acre</b>
Brown Marmorated Stink Bug	2.4 – 3.2	0.019 – 0.025
<b>Notes and Restrictions</b>		
Pre-Harvest Interval (PHI): <b>3 days.</b>		
Maximum BAYTHROID XL allowed per 14-day interval: <b>3.2 fluid oz/A (0.025 lb AI/Acre).</b>		
Maximum BAYTHROID XL allowed per crop season: <b>12.8 fluid oz/A (0.1 lb AI/Acre).</b>		
Minimum application volume (water): 50 GPA – ground, 25 GPA – aerial application.		

**POTATO AND OTHER TUBEROUS AND CORM VEGETABLES**

Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

<b>PEST CONTROLLED</b>	<b>Rate fluid ounces/Acre</b>	<b>Rate lb AI/Acre</b>
Brown Marmorated Stink Bug	1.6 – 2.8	0.013 – 0.022

**Notes and Restrictions**

Pre-Harvest Interval (PHI): **0 day.**

If more than 5.6 fluid oz/Acre is applied, allow at least 14 days between last application and grazing.

Maximum BAYTHROID XL allowed per 5-day interval: **2.8 fluid oz/A (0.022 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **16.8 fluid oz/A (0.132 lb AI/Acre).**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*