

Jun 2, 2011

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration**RESTRICTED USE PESTICIDE**
Due to Toxicity to Fish and Aquatic Organismsin New York State
under 6NYCRR Part 326**For retail sale to and use only by Certified Applicators or persons under their direct supervision
and only for those uses covered by the Certified Applicator's certification.****Bayer CropScience****Bayer CropScience LP**
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina
27709
1-866-99BAYER (1-866-992-2937)**GROUP****3****INSECTICIDE****Baythroid® XL****EPA Reg. No. 264-840****For use on Grape, Potato and other
Tuberous and Corm Vegetables to control
Brown Marmorated Stink Bug**

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

GRAPE

Includes: Table grape, Raisin, Wine and Muscadine grape

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre
Brown Marmorated Stink Bug	2.4 – 3.2	0.019 – 0.025

Notes and Restrictions

Pre-Harvest Interval (PHI): **3 days.**

Maximum BAYTHROID XL allowed per 14-day interval: **3.2 fluid oz/A (0.025 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **12.8 fluid oz/A (0.1 lb AI/Acre).**

Minimum application volume (water): 50 GPA – ground, 25 GPA – aerial application.

POTATO AND OTHER TUBEROUS AND CORM VEGETABLES

Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre
Brown Marmorated Stink Bug	1.6 – 2.8	0.013 – 0.022

Notes and Restrictions

Pre-Harvest Interval (PHI): **0 day.**

If more than 5.6 fluid oz/Acre is applied, allow at least 14 days between last application and grazing.

Maximum BAYTHROID XL allowed per 5-day interval: **2.8 fluid oz/A (0.022 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **16.8 fluid oz/A (0.132 lb AI/Acre).**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)