


# Belmont™ -2.7 FS



Seed Treatment Fungicide  
for control of seed rot and damping-off diseases of certain crops

Active Ingredient:

metalaxyl (CAS No. 57837-19-1) .....	28.98%*
Other Ingredients .....	71.02%
Total .....	100.00%

\*Contains 2.65 lbs. metalaxyl per gallon, or 317 grams per liter.

## KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing</li><li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
24 Hour Emergency Number      1-866-928-0789	
Product Use Information Number 1-866-761-9397	

EPA REG. NO. 400-577  
EPA EST. NO.  
002/121713  
SAP#523873 002 (0115)

Manufactured for:  
MacDermid Agricultural Solutions, Inc.  
245 Freight Street  
Waterbury, CT 06702-1818

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING.** Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes. Do not get on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** Coveralls over short-sleeved shirt and short pants, socks and chemical-resistant footwear and chemical-resistant gloves (such as natural rubber > 14 mils in thickness). When mixing and loading, and when cleaning equipment, wear a chemical-resistant apron. NOTE: Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no worker contact with the soil sub-surface or treated seed. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; socks and chemical-resistant footwear; chemical resistant gloves; and protective eyewear.

## PRODUCT INFORMATION

BELMONT™ 2.7 FS is a systemic fungicide seed treatment specifically for control of systemic downy mildews, Pythium and Phytophthora spp. For control of other soilborne diseases, use in combination with other seed treatment fungicides.

VITAVAX® and RANCONA® products are compatible with BELMONT 2.7 FS. Do not use BELMONT 2.7 FS in combination with other seed treatment products unless compatibility and safety to the crop has been verified. Do not mix with any pesticide that has a prohibition against tank mixing. Observe the most restrictive of labeling limitations and precautions of all products used in mixtures.

BELMONT 2.7 FS may be applied on its own, as water-based slurry, or in combination with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: BELMONT 2.7 FS is a systemic fungicide with a specific mode of action. BELMONT 2.7 FS could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, the company cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible BELMONT 2.7 FS resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

## USE RESTRICTIONS

- Seed treated commercially or for sale must be dyed an un-natural color using an EPA approved colorant before being sold.
- IN CALIFORNIA: ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use in California.
- IMPORTANT: MAXIMUM USAGE when applying multiple metalaxyl and mefenoxam containing products to the same crop within the same season: DO NOT apply more than the maximum seasonal total for the active ingredient as stated on the label of the lowest seasonal total on that crop.
- Do not use other metalaxyl or mefenoxam products for control of Pythium in water-seeded rice.
- Do not apply BELMONT 2.7 FS to spinach which is destined to be grown in the greenhouse.

## TREATED SEED LABELING

Seeds treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This package or bag contains seed that has been treated with metalaxyl. Do not use treated seed for food, feed, oil production or any other purpose except planting. Store away from feed and foodstuffs. Do not allow children, pets or livestock to have access to treated seeds."

"Wear long pants, long-sleeved shirt, shoes and socks, and protective gloves opening bag or handling treated seed."

"Do not use other metalaxyl or mefenoxam products for control of Pythium in water-seeded rice."

"Do not apply metalaxyl to spinach which is destined to be grown in the greenhouse."

"Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess seed by burying away from bodies of water."

"Do not contaminate bodies of water when disposing of planting equipment wash water."

"Dispose of seed packaging or containers in accordance with local requirements."

**APPLICATION INSTRUCTIONS**

<b>Crop</b>	<b>Application Rate Per 100 lbs. of seed</b>	<b>Diseases Controlled</b>
Canola*, Mustard seed*, Rapeseed*  *Not registered for use in CA	0.25 to 0.5 fl. oz.  To deliver 0.00023 - 0.00047 mg.ai. per seed, apply 0.0025 - 0.005 fl. oz. of product per 100,000 seeds (Based on average of 100,000 seeds per lb.)	<i>Pythium</i> damping-off
Cotton	0.75 to 1.5 fl. oz.  To deliver 0.0156 - 0.0312 mg.ai. per seed, apply 0.167 - 0.333 fl. oz. of product per 100,000 seeds (Based on average of 4,500 seeds per lb.)	<i>Pythium</i> seed rot and damping-off
Soybeans	0.75 to 1.5 fl. oz.  To deliver 0.0251 - 0.0502 mg.ai. per seed, apply 0.038 - 0.075 fl. oz. of product per 140,000 seeds (Based on average of 2,800 seeds per lb.)	<i>Pythium</i> damping-off and early season <i>Phytophthora</i>
Peanuts*  *Not registered for use in CA	0.75 fl. oz.  To deliver 0.1004 mg.ai. per seed, apply 1.071 fl. oz. of product per 100,000 seeds (Based on average of 700 seeds per lb.)	<i>Pythium</i> seed rot and damping-off
Rice* *Not registered for use in CA	0.75 to 1.5 fl. oz.  To deliver 0.0037 - 0.0075 mg.ai. per seed, apply 0.040 - 0.080 fl. oz. of product per 100,000 seeds (Based on average of 18,800 seeds per lb.)  Note: Do not use other metalaxyl or mefenoxam containing products in water-seeded rice. Containers or bags containing Belmont 2.7 FS treated rice must be labeled with this restriction	<i>Pythium</i> damping-off
Sorghum / milo Note: Only high quality seed (at least 90% germination) should be treated with BELMONT 2.7 FS.	0.375 to 0.75 fl. oz.  To deliver 0.0020 - 0.0039 mg.ai. per seed, apply 0.021 - 0.042 fl. oz. of product per 100,000 seeds (Based on average of 18,000 seeds per lb.)	<i>Pythium</i> damping-off
	0.75 to 1.5 fl. oz.  To deliver 0.0039 - 0.0078 mg.ai. per seed, apply 0.042 - 0.083 fl. oz. of product per 100,000 seeds (Based on average of 18,000 seeds per lb.)	Systemic downy mildew (races susceptible to chemical control, and cultivars which possess sorghum downy mildew resistance)
	1.5 to 3.0 fl. oz.  To deliver 0.0078 - 0.0156 mg.ai. per seed, apply 0.083 - 0.167 fl. oz. of product per 100,000 seeds (Based on average of 18,000 seeds per lb.)	Systemic downy mildew (races susceptible to chemical control, and cultivars which do not possess sorghum downy mildew resistance)
Sunflowers (oil and confection type)	1.5 to 3.0 fl. oz.  For oil type: To deliver 0.0201 - 0.0402 mg.ai. per seed, apply 0.214 - 0.429 fl. oz. of product per 100,000 seeds (Based on average of 7,000 seeds per lb.)  For confection type: To deliver 0.0402 - 0.0804 mg.ai. per seed, apply 0.429 - 0.857 fl. oz. of product per 100,000 seeds (Based on average of 3,500 seeds per lb.)	Systemic downy mildew
Beans (except soybeans): Field beans, green beans, chick-peas (garbanzos), kidney beans, lima beans, lupines, navy beans, okra, pole beans, snap beans, string beans, wax beans, and lentils.	0.75 fl. oz.  To deliver 0.0439 mg.ai. per seed, apply 0.469 fl. oz. of product per 100,000 seeds (Based on average of 1,600 seeds per lb.)	<i>Pythium</i> seed rot and damping-off and early season <i>Phytophthora</i>

Crop	Application Rate Per 100 lbs. of seed	Diseases Controlled
Peas: Black-eyed peas, cowpeas, field peas, garden peas	<p>0.75 fl. oz.</p> <p>For small size seeds (average of 3,300 seeds per lb.): To deliver 0.0213 mg.ai. per seed, apply 0.227 fl.oz. of product per 100,000 seeds</p> <p>For larger size seeds (average of 2,100 seeds per lb.): To deliver 0.0338 mg.ai. per seed, apply 0.361 fl.oz. of product per 100,000 seeds</p>	<i>Pythium</i> seed rot and damping-off and early season <i>Phytophthora</i>
	<p>1.5 fl. oz.</p> <p>For field peas: To deliver 0.0426 mg.ai. per seed, apply 0.455 fl. oz. of product per 100,000 seeds (Based on average of 3,300 seeds per lb.)</p> <p>For garden peas: To deliver 0.0676 mg.ai. per seed, apply 0.721 fl. oz. of product per 100,000 seeds (Based on average of 2,080 seeds per lb.)</p>	Systemic downy mildew and <i>Pythium</i> damping-off
Corn, sweet	<p>0.75 fl. oz.</p> <p>To deliver 0.0223 mg.ai. per seed, apply 0.238 fl. oz. of product per 100,000 seeds (Based on average of 3,150 seeds per lb.)</p>	<i>Pythium</i> damping-off
	<p>1.5 fl. oz.</p> <p>To deliver 0.0446 mg.ai. per seed, apply 0.476 fl. oz. of product per 100,000 seeds (Based on average of 3,150 seeds per lb.)</p>	Systemic downy mildew
Corn, field	<p>0.75 fl. ozs.</p> <p>To deliver 0.0419 mg.ai. per seed, apply 0.446 fl. oz. of product per 100,000 seeds (Based on average of 1,680 seeds per lb.)</p>	<i>Pythium</i> damping-off
Corn, pop	<p>0.75 fl. ozs.</p> <p>To deliver 0.0230 mg.ai. per seed, apply 0.245 fl. oz. of product per 100,000 seeds (Based on average of 3,060 seeds per lb.)</p>	<i>Pythium</i> damping-off
Cereal Grains: Wheat, barley, oats, rye and triticale	<p>0.75 fl. oz.</p> <p>For wheat, barley, oats: To deliver 0.0054 mg.ai. per seed, apply 0.058 fl. oz. of product per 100,000 seeds (Based on average of 13,000 seeds per lb.)</p> <p>For rye / triticale: To deliver 0.0037 mg.ai. per seed, apply 0.040 fl.oz. of product per 100,000 seeds (Based on average of 18,850 seeds per lb.)</p>	<i>Pythium</i> damping-off
Millet	<p>0.75 fl. oz.</p> <p>To deliver 0.0009 mg.ai. per seed, apply 0.009 fl. oz. of product per 100,000 seeds (Based on average of 80,000 seeds per lb.)</p>	<i>Pythium</i> damping-off
Buckwheat	<p>0.75 fl. oz.</p> <p>To deliver 0.0054 mg.ai. per seed, apply 0.058 fl. oz. of product per 100,000 seeds (Based on average of 13,000 seeds per lb.)</p>	<i>Pythium</i> damping-off

Crop	Application Rate Per 100 lbs. of seed	Diseases Controlled
<p>Cucurbit Vegetables:</p> <p>Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuzza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixtar, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus)</p>	0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Leafy Vegetables:</p> <p>Amaranth (leafy, Chinese spinach**, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (fino- chio), lettuce (head and leaf ), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard.</p> <p>** Do not apply to spinach seed to be grown in greenhouses. Containers or bags containing BELMONT 2.7 FS –treated spinach seed must be labeled with this restriction.</p>	0.1 to 0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Brassica (Cole) Head, Stem and Leaf Vegetables</p> <p>Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens.</p>	0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Forage Legumes:</p> <p>Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).</p>	0.75 to 1.5 fl. oz.	<i>Pythium</i> damping-off and early season <i>Phytophthora</i>
<p>Fruiting Vegetables:</p> <p>Eggplant, groundcherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato</p>	0.1 to 0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Onions (dry bulb and green):</p>	0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Root and Tuber Vegetables (except potatoes)*:</p> <p>Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon),rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manioc pea), yam</p>	0.75 fl. oz.	<i>Pythium</i> damping-off
<p>Turf Grasses:</p> <p>On golf courses, home lawns, reclamation and erosion control</p>	1.5. fl. oz.	<i>Pythium</i> damping-off
<p>Forage Grasses:</p> <p>Grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage</p>	0.75 fl. oz.	<i>Pythium</i> damping-off

**SEED TREATMENT DIRECTIONS:  
USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES**

BELMONT 2.7 FS systemic fungicide seed dressing may be applied at the reduced rates shown in the table below and still provide good Pythium control when used in combination with broad spectrum seed treatment fungicides. Apply as a commercial seed treatment only in combination at EPA registered rates of broad-spectrum seed treatment fungicides for which compatibility and crop safety have been verified. Do not mix with any pesticide that has a prohibition against tank mixing. BELMONT

2.7 FS may be applied on its own or as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

Crop	Application Rate Per 100 lbs. of seed	Diseases Controlled
Cereal Grains Wheat, barley, oats, rye	0.10 to 0.375 fl.oz.  For wheat, barley, oats: To deliver 0.0007 to 0.0027 mg.ai. per seed, apply 0.008 to 0.029 fl.oz. of product per 100,000 seeds (Based on average of 13,000 seeds per lb.)  For rye: To deliver 0.0005 to 0.0019 mg.ai. per seed, apply 0.005 to 0.020 fl.oz. of product per 100,000 seeds (Based on average of 18,850 seeds per lb.)	Seed decay and damping-off caused by Pythium
Corn, field	0.10 to 0.375 fl.oz.  To deliver 0.0056 to 0.0209 mg.ai. per seed, apply 0.060 to 0.223 fl.oz. of product per 100,000 seeds (Based on average of 1,680 seeds per lb.)	Seed decay and damping-off caused by Pythium
Peanuts* *Not registered for use in California	0.20 to 0.375 fl.oz.  To deliver 0.0268 - 0.0502 mg.ai. per seed, apply 0.286 - 0.536 fl.oz. of product per 100,000 seeds (Based on average of 700 seeds per lb.)	Seed decay and damping-off caused by Pythium
Rice* *Not registered for use in California	0.10 to 0.375 fl.oz.  To deliver 0.0005 - 0.0019 mg.ai. per seed, apply 0.005 - 0.020 fl.oz. of product per 100,000 seeds (Based on average of 18,800 seeds per lb.)	Seed decay and damping-off caused by Pythium
Soybeans	0.20 to 0.375 fl.oz.  To deliver 0.0067 - 0.0126 mg.ai. per seed, apply 0.10 - 0.19 fl.oz. of product per 140,000 seeds (Based on average of 2,800 seeds per lb.)	Pythium damping-off and early season Phytophthora
Peas	0.20 to 0.375 fl.oz.  To deliver 0.0059 - 0.0110 mg.ai. per seed, apply 0.063 - 0.117 fl.oz. of product per 100,000 seeds (Based on average of 3,200 seeds per lb.)	Pythium seed rot and damping-off
Beans (Field, kidney, navy, pinto, lentils)	0.20 to 0.375 fl.oz.  To deliver 0.0117 - 0.0220 mg.ai. per seed, apply 0.125 - 0.234 fl.oz. of product per 100,000 seeds (Based on average of 1,600 seeds per lb.)	Pythium seed rot and damping-off
Sunflowers	0.10 to 0.375 fl.oz.  To deliver 0.0017 - 0.0064 mg.ai. per seed, apply 0.018 - 0.068 fl.oz. of product per 100,000 seeds (Based on average of 5,500 seeds per lb.)	Pythium seed rot and damping-off

**SEED TREATMENT DIRECTIONS - FOR EXPORT USE ONLY:**

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation. The treater is responsible for confirming the acceptability of his treated seed in the country to which he is exporting.

<b>Crop</b>	<b>Application Rate Per 100 lbs. of seed</b>	<b>Diseases Controlled</b>
Brassica (Cole) Head, Stem & LeafyVegetables	0.75 - 1.5 fl. oz.	Pythium damping-off
Canola	0.5 – 1.70 fl. oz. (32 to 110 ml per 100 kg. of seed)	Pythium damping-off
Corn	0.75 - 1.5 fl. oz.	Pythium damping-off
	3.0 fl. oz.	Systemic downy mildew and Pythium damping-off
Cucurbit Vegetables	0.75 fl. oz.	Pythium damping-off
Fruiting Vegetables	0.75 - 1.5 fl. oz.	Pythium damping-off
Leafy Vegetables	0.75 - 1.5 fl. oz.	Pythium damping-off
Millet	0.75 - 1.5 fl. oz.	Pythium damping-off
	2.5 fl. oz.	Systemic downy mildew and Pythium damping-off
Onions	0.75 - 1.5 fl. oz.	Pythium damping-off
Peas	3.5 fl. oz.	Pythium damping-off and early season Phytophthora
Root and Tuber Vegetables (except potatoes)	0.75 - 1.5 fl. oz.	Pythium damping-off
Sorghum	0.75 - 1.5 fl. oz.	Pythium damping-off
	5.25 fl. oz.	Systemic downy mildew and Pythium damping-off
Sunflowers	9.0 fl. oz.	Systemic downy mildew

**TREATED SEED LABELING:**

Seed treated in accordance with the Export Use pattern must be labeled as follows:  
“FOR EXPORT ONLY – NOT FOR DOMESTIC SALES OR USE”.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal or by cleaning of equipment.

**PESTICIDE STORAGE:** Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Non-refillable container

Plastic Containers: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Triple rinse as follows:** *For containers with capacity equal to or less than 5 gallons or 50 pounds:* Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume - and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

*For containers with capacities greater than 5 gallons or 50 pounds:* Empty the remaining contents into application equipment or a mix tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

**Recycling:** Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or [www.acrecycle.org](http://www.acrecycle.org).

**Refillable container**

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**To clean the container before final disposal,** empty the remaining contents into application equipment or mix tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

## IMPORTANT: READ BEFORE USE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond control of MacDermid Agricultural Solutions, Inc., MacDermid Agricultural Solutions, Inc. makes no claims of guarantees as to germination of carry-over seed.

BELMONT™, VITAVAX® and RANCONA® are trademarks of MacDermid Agricultural Solutions, Inc.

©Copyright 2015, MacDermid Agricultural Solutions, Inc.