

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



EPA Reg No. 71711-61

For Use on Citrus Fruit (Crop Group 10-10)

States: California Only

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for **BEXAR CA** Insecticide before applying. Use of **BEXAR CA** Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **BEXAR CA** Insecticide.

APPLICATION RATE CHART FOR BEXAR CA INSECTICIDE

Citrus Fruit (Crop Group 10-10) Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin orange, clementine); tangor; trifoliate orange; unqi fruit; cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Glassy-winged sharpshooter	21.0 to 27.0 fl oz/acre	<ul style="list-style-type: none"> • Apply only to groves practicing weed management. • Apply by ground only, using a minimum of 100 gallons of water per acre. • For low volume or air curtain sprayers, apply using a minimum of 30 gallons of water per acre. <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Do not apply by air on citrus fruit. • Do not spray during blooming period or when bees are foraging in the field. • Do not apply this product or allow it to drift to blooming crops, plants, or weeds while bees are foraging in or adjacent to the treatment area. • If bees are foraging in vegetative filter strips that contain any blooming plants or weeds, always remove flowers before making an application. This may be accomplished by mowing or applying a labeled herbicide. • Do not use on nursery stock. • Do not apply to the first row of outer perimeter trees and to trees at each end of the tree row. • Do not apply more than 27.0 fl oz per acre per growing season. • Do not make more than 1 application every other calendar year. • Preharvest Interval (PHI): 3 days

©2021 Nichino America, Inc. Bexar and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700