



BIO-ACTIVATE

Natural Plant Growth Factor



INGREDIENTS

Extract of microbial culture.....	1.1%
Mixture of enzymes and amino acids.....	1.0%
Growth stimulating factors.....	0.1%
(cytokinin activity)	
Inert Ingredient.....	98.9%

DIRECTIONS FOR USE

BIO-ACTIVATE should be applied with enough water to thoroughly cover all plants. Apply **BIO-ACTIVATE** early in the morning or late in the afternoon.

RATE OF APPLICATION

16 - 24 oz. of **BIO-ACTIVATE** per hectare.

TIME OF APPLICATION

Apple: Apply at early spring when new leaves are emerged and at early bloom.

Carrot: Apply prior to tuber initiation.

Celery: Apply at 2-3 leaf stage.

Citrus: Apply at early spring when new leaves are emerged and at early bloom.

Peach: Apply at pit-hardening stage.

Potato First application should be made when crops are about 4 to 6 inches. Repeat the application in 2-3 weeks.

Wheat: Apply at 4-6 leaf stage.

BIO-ACTIVATE is a natural plant growth factor derived from microbial cultures. It contains a mixture of cytokinins which are known to regulate plant growth.

NO WARRANTY

Neither manufacturer nor sellers shall be liable for any direct or indirect damage resulting from the use or handling of this product whether or not in accordance with directions or suggestions. The user assumes all risks in use or handling. Since weather, crop, soil and other conditions may vary, the manufacturer and the seller make no warranty concerning the effectiveness of the product.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Net Contents:

- ☐ 1 Pint (0.47 liter)
 - ☐ 2.5 Gallons (9.45 liters)
 - ☐ 2 x 2.5 Gallons (18.9 liters)
 - ☐ 5 Gallons (18.9 liters)
 - ☐ 55 Gallons (208 liters)
 - ☐ Other: _____ Gallons
- Net Wt. 9.0 lbs. per gallon

Lot No.:

Manufacture Date:

JH Biotech Inc.

0221-D-A-1
Copyright © JH Biotech, Inc.
All right reserved.