

Bio-Wet

Wetting Agent and Humectant

DIRECTIONS FOR USE: SHAKE WELL Before use.

The effectiveness of pesticides and foliar feeds may be significantly impacted by the wetting and retention properties of the spray solution. Better retention provides spray solutions with a longer window of time in which to work, thereby increasing their effectiveness.

Bio-Wet can be used to ensure optimum results by improving the retention, wetting, spreading and adhesion of the pesticidal or fertilizer deposits.

CAUTION: Bio-Wet must be thoroughly mixed with water before adding pesticides or foliar feeds. The rate of Bio-Wet must be adjusted according to the volume of the application (ground, concentrate, aerial) and the plant surface to be sprayed (waxy, hairy). This product is prohibited from use on aquatic sites.

WA Reg. No. 99891-10001

Batch No. 221209

A soluble wetter and humectant for use in certified organic product production to improve retention and adhesion of pesticides and foliar feeds.

Principal Functioning Agents: Yucca, Saccharum and Ascophyllum nodosum extract	92.5%
Constituents Ineffective as Spray Adjuvants	7.5%
Total	100%

- Bio-Wet contains a combination of nonionic and anionic components
- Bio-Wet ensures adequate distribution of spray material over difficult to wet, waxy plant surfaces.
- Bio-Wet decreases unnecessarily high spray run-off.
- Bio-Wet reduces the possibility of chemical damage to plants by evenly distributing spray materials.



The following table serves as an indicator of the quantity of Bio-Wet required for various application types.

AMOUNT OF BIO-WET TO BE ADDED TO VARIOUS APPLICATION TYPES		
APPLICATION TYPE	fl. oz/100 gallons of water	mL/100 liters of water
High volume ground spraying (218 – 428 gal water/acre) (2000 – 4000 L water/ha)	1.9 – 2.6 fl. oz	15 – 20 mL
Concentrate spraying (low volume) (54-107 gal water/acre) (500-1000 L water/ha)	2.6 – 4.5 fl. oz	20 – 35 mL
Conventional ground spraying (21-54 gal water/acre) (200-500 L water/ha)	4.5 fl. oz	35 mL
Aerial spraying (2.7-4.8 gal water /acre) (25-45 L water/ha)	12.5 fl. oz	100 mL

N.B. Always mix Bio-Wet thoroughly with water before adding pesticides or foliar feeds.

WARRANTY: The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences to the environment, wildlife or humans may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of NutriAg Ltd. To the extent consistent with applicable law, NutriAg Ltd. shall not be liable for consequential, financial, special or indirect damages resulting from the use or handling of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer. The seller warrants that the product conforms to its chemical description. Consult the supplier in any event of uncertainty.

CAUTION

PRECAUTIONARY STATEMENTS: Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT: Wear protective eyewear (goggles or face shield), chemical-resistant gloves, long sleeved shirt and long pants, and shoes plus socks.

FIRST AID:

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911, or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

STORAGE AND DISPOSAL: Store in original container only. Do not store in metal container. Keep away from strong oxidizing agents and acids. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option for container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler, visit the ACRC web page at www.acrcycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

Net Contents: 2.5 gal. (9.46 L)
Net Weight: 26.08 lbs. (11.83 kg)

NutriAg Ltd, 39 Gail Grove, Toronto ON, Canada, M9M 1M5 Tel: 416-636-1555