



BioFlora® Boron-Zinc (1%-4%)

BioFlora® Boron-Zinc (1%-4%) is designed to strengthen cell walls and aids in carbohydrate, protein, and auxin metabolism. It aids in the correction of boron and zinc deficiencies. It is a co-factor for various enzymes taking part in cellular processes. Best results are achieved when applications are made to actively growing crops after an irrigation. This product is enhanced with our patented process known as Cellular Infuson® Technology. Driven by a nanomolecule and ionic minerals, Cellular Infuson® Technology improves the absorption, bioavailability and translocation of nutrients in plants at the cellular level.

DISCLAIMERS:

This product does not contain microorganisms.

KEEP AWAY FROM CHILDREN. NOT FOR HUMAN CONSUMPTION.

Buyer assumes all risk of use, storage and handling of this material. Neither seller nor its agent make any warranty, expressed or implied, concerning this product, except in conformity with statement on the label.

Information regarding the contents and levels of metals in this product is available online at aapfco.org/metals.

STORAGE & HANDLING:

Shake well before use. Store in original container in a cool dry place out of direct sunlight. In case of accidental exposure, flush with water. Wear eye protection. Protective gloves recommended.



www.BioFlora.com

info@BioFlora.com

(623) 932-1522

GUARANTEED ANALYSIS:

Boron (B)1.00%
Zinc (Zn)4.00%

DERIVATION:

Derived from disodium octaborate and zinc sulfate

DENSITY:

9.40 lb per gallon
1.18 kg per liter

COMPATIBILITY:

Do not mix with products containing materials with a high pH.

DIRECTIONS FOR USE:

To achieve thorough coverage use adequate water. For trees apply 1 gallon in 500 gallons of dilute spray. For air applications use 5-10 gallons of water/acre.

When used with other materials the mixing sequence in the spray tank will be: water, **BioFlora® Boron-Zinc**, fertilizer or pesticide. Apply at temperatures below 90°F (32°C).

APPLICATION RATES:

Foliar: apply 1 - 2 quarts of concentrate per acre every 2 weeks. Up to 2 gallons per acre for a growing season may be used. Product should be mixed with 40 - 50 gallons of water prior to application.

