

BioFlora® Zinc

FOR AGRICULTURE

5 Gallons (18.93 liters) • Net Weight 53.50 lb (24.23 kg)

LOT # 006445

GUARANTEED ANALYSIS:

Zinc (Zn) 7.00%

DERIVATION:

Derived from zinc sulfate.

DENSITY:

10.70 lb per gallon; 1.28 kg per liter

DIRECTIONS FOR USE:

For correction of zinc deficiencies of field, row, vegetable, fruit, nut,, vine, ornamental and turf crops. **BioFlora® Zinc**, alone, or with pesticides or fertilizers, may be applied directly to soil or to the foliage of growing crops. **BioFlora® Zinc** may be applied via ground sprayers, aircraft or drip irrigation systems. Shake, stir or swirl contents before using. **MIXING: Always add this product to spray tank before adding pesticides.**

COMPATIBILITY:

BioFlora® Zinc is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

KEEP AWAY FROM CHILDREN. NOT FOR HUMAN CONSUMPTION.

Note: Buyer assumes all risks of use, storage and handling of this material. Neither seller nor its agent make any warranty, expressed or implied, concerning this product, except in conformity with statements on the label.

DISCLAIMER: This product does not contain microorganisms.

BIOFLORA SYSTEMS
Manufactured & Guaranteed by:
GLOBAL ORGANICS, LLC
16121 W. EDDIE ALBERT WAY
GOODYEAR, AZ 85338
TELEPHONE: 623-932-1522 WEB: www.bioflora.com

APPLICATION RATES:

Soil applications of **BioFlora® Zinc** are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Immediate post-irrigation or post-rainfall applications give best foliar uptake.

SOIL APPLICATIONS:

Soil Fertility Maintenance 1 - 3 qts/acre
Mild Deficiency 1 - 3 qts/acre
Severe Deficiency 2 - 6 qts/acre

*****ALL CROPS:** The lower rates are for field, row and vegetable crops. The higher rates are for fruit, nut, vine, ornamental and turf crops.

FOLIAR APPLICATIONS: Apply 2 - 4 qts/acre as needed. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

DILUTE APPLICATIONS: Apply the suggested dose of product in enough water for good coverage.

CONCENTRATE APPLICATIONS: Use up to 2 quarts in a minimum of 25 gallons of water.

AERIAL APPLICATIONS: Up to 1 quart per gallon of spray may be used, but more dilute solutions will improve coverage and results.

DRIP IRRIGATION APPLICATIONS: Apply the recommended amount of product in split applications 1 - 2 weeks apart throughout the growing season.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

