

POWDER

BIOMIN[®] CALCIUM

Chelated Minerals for Soil and Foliar Application

5-0-0

Product Information

BIOMIN CALCIUM contains nitrogen and chelated calcium for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. **BIOMIN CALCIUM** is designed to prevent and correct calcium deficiency and to boost crops during critical or fast growing periods. **BIOMIN CALCIUM** may be applied with nitrogen fertilizers.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	5.0%
5% Water Soluble Organic Nitrogen	
Calcium (Ca).....	15.0%
15% Chelated Calcium	

Derived from calcium carbonate, citric acid (chelating agent), and soy protein hydrolysate.

KEEP OUT OF REACH OF CHILDREN

COMPATIBILITY: Always perform a compatibility test before mixing any products with **BIOMIN CALCIUM**.

PREVENTS AND CORRECTS CALCIUM DEFICIENCY

NET WEIGHT: 25 lb. (11.3 kg)
U.S. Pat.# 5,504,055
Lot #:



MANUFACTURED BY: JH BIOTECH, INC.

P.O. Box 3538, Ventura, CA 93006 U.S.A.

805-650-8933 jhbiotech.com

Directions and General Recommendations

SOIL APPLICATION

BIOMIN CALCIUM may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5-5.0 lb./ac. (1.7-5.6 kg/ha) depending on crop, soil conditions and application methods. For maintenance, 1.5 lb./ac. (1.7 kg/ha) are usually recommended.

FOLIAR APPLICATION

FIELD AND ROW CROPS: Apply 1.0-1.5 lb./ac. (1.1-1.7 Kg/Ha) of **BIOMIN CALCIUM** when crops are 6-12 in. (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

VEGETABLES AND POTATOES: Apply 1.0-1.5 lb./ac. (1.1-1.7 kg/ha) of **BIOMIN CALCIUM** at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

GREENHOUSE CROPS: Apply 1.0-1.5 lb./ac. (1.1-1.7 kg/ha) of **BIOMIN CALCIUM** 2-3 weeks after emergence or transplanting, or when calcium deficiencies first appear. Repeat the application every one or two weeks.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

FRUIT TREES AND GRAPES: Apply 1.0-1.5 lb./ac. (1.1-1.7 kg/ha) of **BIOMIN CALCIUM** in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

BIOMIN CALCIUM is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, dilute it with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with **BIOMIN CALCIUM**.

BIOMIN CALCIUM can be applied to all commercial crops including all vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers and pastures.

Precautionary statement: do not swallow. Wash hands thoroughly after handling. **First Aid:** if swallowed: Seek medical attention if feeling unwell. Rinse mouth. **Storage & Disposal:** store in a safe manner. Dispose product/container in accordance with federal, state, and local regulations.

052003-D-X-1

Copyright © 2017 All rights reserved