

# Biscayne™



### Agricultural, Commercial Noncropland Sites and Turf and

**Ornamental Uses:** This product may be used as a postemergence application to control sedges and broadleaf weeds in peanuts, corn, beans, clover grown for seed (Washington and Oregon Only), peas, rice, soybeans, sorghum, spearmint and peppermint. This product may also be used to control annual sedges, broadleaf weeds and yellow nutsedge in ornamentals, turfgrass and other noncropland sites listed in the Directions for Use.

<b>ACTIVE INGREDIENT:</b>	(% by weight)
Sodium salt of bentazon* [sodium 3-(1-methylethyl)-1H-2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide] .....	44.0%
<b>OTHER INGREDIENTS:</b> .....	56.0%
<b>TOTAL:</b> .....	100.0%

\* Equivalent to 4 pounds of bentazon per gallon.

EPA Reg. No.: 91234-102

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

### FIRST AID

**If on skin or clothing:** ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice. **If swallowed:** ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person. **If in eyes:** ▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,  
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long sleeved shirt and long pants, shoes plus socks and chemical-resistant gloves. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are made of waterproof material.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Bentazon, which is present in this product, is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Notice:** It is a violation of federal law to use any pesticide in a manner that results in the death of an endangered species or in adverse modification of their habitat.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are to be followed. This labeling must be in the user's possession during application.

### Pollinator Advisory Statement

This product may adversely impact the forage and habitat of local pollinators, including the monarch butterfly (and its larvae), birds, or bats if reaches non-target areas. Protect pollinators by following label directions to minimize spray drift.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

**For Non-WPS Occupational Use:** Do not enter or allow others to enter the treated area until sprays have dried.

### WEED RESISTANCE MANAGEMENT

For resistance management, **Biscayne** is a Group 6 herbicide. Any weed population may contain or develop plants naturally resistant to **Biscayne** and other Group 6 herbicides. Weed species with acquired resistance to Group 6 herbicides may eventually dominate the weed population if Group 6 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **Biscayne** or other Group 6 herbicides. Users should scout before and after application.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance:

- Avoid the consecutive use of **Biscayne** or other target site of action Group 6 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern (an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides).

- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Report any incidence of non-performance of this product against a particular weed species to your Atticus, LLC retailer or representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

### SPRAY DRIFT MANAGEMENT

Bentazon can affect non-target plant species outside the treatment area. To limit adverse effects to non-target plants, the applicator must avoid making applications when wind can facilitate off-site movement of bentazon in the direction of areas such as forested areas, riparian areas, wetlands, and areas that serve as habitat for desirable and protected animal species. **DO NOT** apply by air if sensitive crop species (such as cotton, sugar beets, sunflowers, or okra) are within 200 feet downwind.

#### SPRAY DRIFT

##### Aerial Applications:

- When applying aerially to crops, do not release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver medium or coarser spray droplets in accordance with ASABE Standard S-572.1.
- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

##### Ground Boom Applications:

- When using ground application equipment, apply with nozzle height no more than 4 feet above the ground or crop canopy.
- Applicators are required to select nozzles that deliver medium or coarser spray droplets in accordance with ASABE Standard S-572.1.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

### SPRAY DRIFT ADVISORIES

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient

coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **WIND, TEMPERATURE AND HUMIDITY**, and **TEMPERATURE INVERSIONS** sections of this label.

### Controlling Droplet Size – Ground Boom

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

### Controlling Droplet Size – Aircraft

- Number of Nozzles - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations. **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**
- Nozzle Type - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Boom Length - Longer booms increase drift potential. Therefore a shorter boom length is recommended.
- Application Height - Application more than 10 feet above the canopy increases the potential for spray drift.

### BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator needs to be familiar be familiar with local wind patterns and how they affect spray drift.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally

in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

## RUNOFF PREVENTION

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

## AGRICULTURAL USES

### PRODUCT INFORMATION

**Biscayne** is a selective herbicide for postemergence control of listed sedges and broad-leaf weeds in beans, peanuts, corn, clover grown for seed (Washington and Oregon only), peas, rice, soybeans, sorghum, spearmint and peppermint.

As **Biscayne** controls through contact with the target weed, thorough coverage of weeds with spray solution is essential for an effective application.

All crops listed in this label are tolerant to **Biscayne**. Treatment of listed crops may cause bronzing or leaf speckling, although crops will usually recover within 10 days. Crops will develop normally and crop vigor will not be affected.

Application equipment must be thoroughly cleaned before and after applying **Biscayne** as follows:

1. Use a commercial spray equipment cleaner or a strong detergent in accordance with the manufacturer's directions.
2. Triple rinse application equipment prior to and after application.

### MODE OF ACTION

Bentazon, the active ingredient in **Biscayne**, is a Group 6 (WSSA) herbicide belonging to the benzothiadiazinone chemistry class. **Biscayne** inhibits photosynthesis at photosystem II site B resulting in symptoms of chlorosis that progresses to necrosis and control of emerged weeds.

### APPLICATION INSTRUCTIONS

**Biscayne** may be applied as a spot spray application, a banded application, or as a broadcast application to actively growing weeds. Application rates and growth stages are listed below. **Biscayne** is most effective in controlling target species when it is applied as a postemergence treatment when weeds are young. Applying this product at an early stage provides the most effective treatment (except for Canada thistle and yellow nutsedge) as it allows use of the lower listed application rates (as appropriate to the target species) and spray coverage is easier to achieve. Target species must be covered thoroughly with **Biscayne**. Smaller weeds may shelter under dense leaf canopies, preventing sufficient spray coverage.

Apply **Biscayne** at the rates specified below (as appropriate to the crop site) to target species that are actively growing but before they reach the listed maximum stage of growth. Delayed treatment allows target species to exceed the listed growth stage for application which will limit control.

If the wind speed is greater than 10 mph or conditions promote spray drift from the application site, do NOT apply this product.

## Irrigated Areas

In areas that are irrigated, irrigation prior to treatment may be necessary in order to ensure target species are actively growing. Treatment of weeds growing under conditions of drought may lead to limited control.

## Cultivation

Do not cultivate areas to be treated within 5 days prior to making an application with **Biscayne**.

Do not cultivate treated areas for 7 days following treatment. Cultivation shortly after the 7 day period following application may assist in providing control for the season.

## AERIAL APPLICATION

Apply **Biscayne** using spray equipment with diaphragm-type nozzles producing a fan or cone spray pattern, at a pressure of up to 40 psi, in 5 gallons of water (minimum) per acre. When applying this product to rice, use a minimum of 10 gallons of water per acre.

Spray nozzles must:

1. Be directed so that they discharge straight back with the air stream or at some angle between straight back and straight down.
2. Be within 10 feet above the crop.

## Special Directions for Aerial Application

To obtain uniform coverage and to avoid drift hazards, follow these guidelines:

- Do not apply this product by aircraft when wind is blowing more than 10 mph (except above 5 mph in California).
- Use coarse sprays (larger droplets) as they are less likely to drift.
- Do not apply this product by air if sensitive species (such as cotton, sugar beets, sunflowers, or okra) are within 200 feet downwind.

The applicator must follow the most restrictive use precautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

## GROUND BROADCAST APPLICATION

Apply **Biscayne** using high pressure spray application equipment with hollow cone or flat fan nozzles spaced at a maximum of 20 inches apart at a minimum of 40 psi. Note: Measure spray pressure at the boom, not in the line or at the pump. For the most effective application, use 10 - 20 gallons of spray solution per broadcast acre. If **Biscayne** is applied in areas where weed foliage is thick or when using the lower volume of spray solution (10 gallons per broadcast acre), use a spray pressure of 60 psi (minimum).

## Special Directions for Ground Application

- Do not apply this product using whirl chamber, controlled droplet indicator or flood nozzles as they may cause erratic coverage, which may lead to inconsistent weed control.
- Selective application equipment must not be used as good coverage must be achieved for the most effective results (e.g. wiper applications or recirculating sprayers).

**Biscayne** can be used for control of listed weed species in the following crops:

- Corn
- Dry Peas
- Rice
- Spearmint
- Clover Grown for Seed
- Peanuts
- Sorghum
- Succulent Beans
- Peppermint
- Soybeans
- Succulent Peas
- Dry Beans

## Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice

IMPORTANT: Refer to **CROP SPECIFIC DIRECTIONS** for **Restrictions** and **Precautions** specific to the treated crop.

Weeds Controlled	Application Rate (Pints)	Growth Stage (Leaves)	Max. Height (Inches)	Comment
Anoda, Spurred	1.0	Not Rec.	—	
	1.5	1 - 6	3	
	2.0	6 - 8	4	
Balloonvine	1.0	Not Rec.	—	
	1.5	2 - 4	2	
	2.0	4 - 6	3	
Beggarticks	1.0	Not Rec.	—	
	1.5	1 - 6	6	
	2.0	6 - 8	8	
Bindweed (Field, Hedge)	1.0	Not Rec.	—	For suppression only, apply 2 - 3 pints per acre in IN, IL, KY, MI and OH.
	1.5	Not Rec.	—	
	2.0	—	10	
Buckwheat, Wild	1.0	Not Rec.	—	
	1.5	1 - 4	3	
	2.0	4 - 6	5	
Canada Thistle	1.0	Not Rec.	—	Apply same rate 7 - 10 days later if regrowth occurs.
	1.5	Not Rec.	—	
	2.0	—	8 to bud	
Cocklebur	1.0	2 - 4	4	Do not apply <b>Biscayne</b> before the specified leaf stage. Do not count cotyledon leaves. In order to make a late rescue application for Cocklebur, make one treatment with <b>Biscayne</b> at a rate of 2 - 3 pints per acre to Cocklebur up to 24 inches in height. For control, apply <b>Biscayne</b> at a rate of 1.5 pints per acre. Make another application at the same rate 10 - 14 days later.
	1.5	2 - 6	6	
	2.0	6 - 10	10	
Croton, Tropic	1.0	Not Rec.	—	
	1.5	up to 2	2	
	2.0	2 - 4	4	
Dayflower	1.0	Not Rec.	—	
	1.5	up to 6	4	
	2.0	6 - 10	8	
Devilsclaw	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	Not Rec.	—	
	2.0	up to 6	3	

(continued)

## Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice (continued)

IMPORTANT: Refer to **CROP SPECIFIC DIRECTIONS** for **Restrictions** and **Precautions** specific to the treated crop.

Weeds Controlled	Application Rate (Pints)	Growth Stage (Leaves)	Max. Height (Inches)	Comment
Eclipta	1.0	—	—	
	1.5	up to 6	2	
	2.0	up to 6	2	
Galinsoga	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	Not Rec.	—	
	2.0	Cotyledon to 6	2	
Groundsel, Common	1.0	Not Rec.	—	
	1.5	Not Rec.	—	
	2.0	—	3	
Jimsonweed	1.0	up to 4	4	
	1.5	up to 6	6	
	2.0	6 - 10	10	
Ladysthumb	1.0	up to 4	4	
	1.5	up to 6	6	
	2.0	6 - 10	10	
Lambsquarters, Common	1.0	up to 4	1	Use crop oil concentrate plus UAN or just crop oil concentrate. If new germination or regrowth occurs, a second treatment with <b>Biscayne</b> may be required.
	1.5	up to 6	1.5	
	2.0	up to 6	2	
Marshelder	1.0	Not Rec.	—	
	1.5	up to 4	2	
	2.0	up to 8	4	
Mayweed/Dogfennel	1.0	Not Rec.	—	
	1.5	—	2	
	2.0	—	3	
Morningglory	1.0	Not Rec.	—	
	1.5	4	4	
	2.0	6	6	

(continued)

### Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice (continued)

IMPORTANT: Refer to **CROP SPECIFIC DIRECTIONS** for **Restrictions** and **Precautions** specific to the treated crop.

Weeds Controlled	Application Rate (Pints)	Growth Stage (Leaves)	Max. Height (Inches)	Comment
Morningglory (Smallflower, Cypressvine Only)	1.0	Not Rec.	—	Rates given for AL, AR, FL, GA, LA, MS, NC, OK, SC, TN, TX, and VA only. Apply a second treatment 7 - 14 days later. For all other states, apply <b>Biscayne</b> at a rate of 2 - 3 pints per acre to annual morningglories up to the 4 true leaves stage of growth. Control of target weed species may be inconsistent or partial.
	1.5	4	4	
	2.0	4	4	
Mustard, Wild	1.0	up to 4	2	
	1.5	up to 6	4	
	2.0	6 - 10	8	
Nightshade, Hairy	1.0	Not Rec.	—	<b>Biscayne</b> does not provide control of black nightshade or Eastern black nightshade.
	1.5	Not Rec.	—	
	2.0	2 - 6	4	
Nutsedge, Yellow	1.0	Not Rec.	—	If regrowth of target species occurs, apply <b>Biscayne</b> a second time at the same rate 7 - 10 days later.
	1.5	—	8	
	2.0	—	8	
Poinsettia, Wild	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	up to 6	4	
	2.0	4 - 8	6	
Purslane, Common	1.0	Not Rec.	—	
	1.5	up to 4	1	
	2.0	4 - 6	2	
Radish, Volunteer	1.0	Not Rec.	—	
	1.5	2 - 6	4	
	2.0	6 - 10	10	
Ragweed, Common	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	Not Rec.	—	
	2.0	4 - 6	3	
Ragweed, Giant	1.0	Not Rec.	—	If new germination or regrowth occurs, a second treatment with <b>Biscayne</b> may be required.
	1.5	Not Rec.	—	
	2.0	up to 4	6	

(continued)

### Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice (continued)

IMPORTANT: Refer to **CROP SPECIFIC DIRECTIONS** for **Restrictions** and **Precautions** specific to the treated crop.

Weeds Controlled	Application Rate (Pints)	Growth Stage (Leaves)	Max. Height (Inches)	Comment
Redweed	1.0	Not Rec.	—	
	1.5	4 - 6	6	
	2.0	6 - 10	8	
Senna, Coffee	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	Not Rec.	—	
	2.0	up to 1 pinnate	2	
Sesbania	1.0	Not Rec.	—	Use crop oil concentrate plus UAN or just crop oil concentrate.
	1.5	Not Rec.	—	
	2.0	3 - 5	3	
Shepherdspurse	1.0	Not Rec.	—	Do not apply to the treat rosette before the seed stalk emerges.
	1.5	up to 6	4	
	2.0	6 - 10	8	
Sida, Prickly or Teaweed	1.0	Not Rec.	—	
	1.5	up to 6	3	
	2.0	6 - 8	4	
Smartweed, Pennsylvania	1.0	up to 4	4	
	1.5	up to 6	6	
	2.0	6 - 10	10	
Starbur, Bristly	1.0	Not Rec.	—	
	1.5	up to 4	2	
	2.0	4 - 6	3	
Sugar Beet, Volunteer	1.0	Not Rec.	—	
	1.5	2 - 4	—	
	2.0	4 - 8	—	
Sunflower, Wild	1.0	up to 2	3	
	1.5	up to 4	5	
	2.0	4 - 6	8	

(continued)

## Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice (continued)

IMPORTANT: Refer to **CROP SPECIFIC DIRECTIONS** for **Restrictions** and **Precautions** specific to the treated crop.

Weeds Controlled	Application Rate (Pints)	Growth Stage (Leaves)	Max. Height (Inches)	Comment
Velvetleaf	1.0	up to 4	2	In order to make a late rescue application for Velvetleaf, make one treatment with <b>Biscayne</b> at a rate of 3 pints per acre with 1 gallon of UAN solution and 1 quart of oil concentrate per acre to velvetleaf plants that are up to 12 inches in height. For control, apply <b>Biscayne</b> at a rate of 1.5 pints per acre plus 1 gallon of AMS or UAN solution and 1 quart of oil concentrate per acre. Then make one more application at the same rate 7 days later. Applicators must use AMS or UAN as a spray additive.
	1.5	up to 4	2	
	2.0	4 - 6	5	
Venice Mallow	1.0	up to 4	2	
	1.5	up to 6	2	
	2.0	6 - 10	4	

### ADDITIVES

In order to achieve control of target weed species on a consistent basis, use one of the following additives:

- Crop Oil Concentrate
- Urea Ammonium Nitrate
- Ammonium Sulfate

The use of additives may cause leaf burn. Leaf burn is more likely when temperature and relative humidity are high. However, crop vigor will not be reduced, and crop growth will be normal. Refer to the **Additive Rates** table below for specific rate instructions.

### Oil Concentrate

Appropriate oil concentrate must either contain a vegetable oil base or a petroleum base. It must also contain EPA-exempt ingredients only and it must not be phytotoxic. The oil concentrate used must have been successful in the locale, and it must display good mixing properties in a jar test.

Although the content of appropriate additive products will differ, petroleum based and vegetable oil based additive products should have emulsifiers in them which will provide good mixing properties. Vegetable oils that are highly refined have been shown to be better for this purpose than vegetable oils that are unrefined.

Refer to the **MIXING INFORMATION** section for more information.

The use of oil concentrate may cause leaf burn. Leaf burn is more likely when temperature and relative humidity are high. However, crop vigor will not be reduced, and crop growth will be normal.

Certain oil concentrate additives can cause excessive leaf burn. Refer to a local supplier for information on the success of the additive in the local area prior to purchase.

### Oil Concentrate plus Nitrogen Solution

Applicators may add an oil concentrate that is nonphytotoxic with a nitrogen solution (i.e. AMS or UAN) to the **Biscayne** spray solution.

### UAN (Urea Ammonium Nitrate)

UAN is often referred to as 28%, 30% or 32% nitrogen solution. UAN can be used instead of other additives in order to achieve control of devilsclaw, cocklebur, velvetleaf, Pennsylvania smartweed, wild mustard, wild sunflower and Venice mallow.

If common lambsquarters and/or common ragweed are present in addition to velvetleaf, (or other weed species that require the use of an oil concentrate) use an oil concentrate. This product combined with a nitrogen solution will not adequately control common lambsquarters and common ragweed.

### AMS (Ammonium Sulfate)

When using AMS, combine 3 quarts of liquid AMS (8-8-0 analysis) with the spray solution or 2.5 pounds of granular AMS. Applicators must only use fine spray grade or feed-grade AMS. AMS that is of an inferior grade will not dissolve fully and may plug spray equipment.

Apply AMS in 10 gallons spray solution per acre minimum. Application of AMS in less than the minimum spray volume may cause problems with precipitation in reduced volumes of water.

Only use AMS if it has been proved to be successful in local area.

### Additive Rates

Additive	Ground Application Rate Per Acre	Air Application Rate Per Acre
UAN Solution*	4 - 8 pints	2 - 4 pints
AMS*	2.5 pounds	2.5 pounds**
Oil Concentrate	1 - 2 pints	1 pint
Oil Concentrate + Nitrogen*	0.5 - 1 pint of Oil Concentrate + 1 - 2 pounds of AMS or 2 - 4 pints of UAN	

\* AMS and UAN must not be used in California.

\*\* Use of AMS solution is not recommended because of precipitation problems in reduced water volumes. Only use AMS when the source has been proved to be successful in the local area and when applied in 10 gallons of solution per acre (minimum).

## MIXING INFORMATION

The following registered products and/or additives may be mixed with **Biscayne**:

Product (Active Ingredient)	Product (Active Ingredient)
<b>Atrazine</b>	<b>Paramount</b> <sup>®</sup> (quinclorac)
<b>Atrazine + dicamba</b>	<b>Paraquat</b>
<b>Avalon</b> <sup>™</sup> (acifluorfen)	<b>Poast</b> <sup>®</sup> (sethoxydim)
<b>Buctril</b> <sup>®</sup> (bromoxynil)	<b>Poast Plus</b> <sup>®</sup> (sethoxydim)
<b>Chlorimuron + Thifensulfuron</b>	<b>Propanil</b>
<b>Clarity</b> <sup>®</sup> (dicamba)	<b>Pursuit</b> <sup>®</sup> (imazethapyr ammonium)
<b>Classic</b> <sup>®</sup> (chlorimuron-ethyl)	<b>Raptor</b> <sup>®</sup> (imazamox ammonium)
<b>Cobra</b> <sup>®</sup> (lactofen)	<b>Reflex</b> <sup>®</sup> (fomesafen)
<b>Concert</b> <sup>®</sup> II (thifensulfuron + chlorimuron-ethyl)	<b>Resource</b> <sup>®</sup> (flumiclorac)
<b>Distinct</b> <sup>®</sup> (diflufenzopyr + dicamba)	<b>Scepter</b> <sup>®</sup> (imazaquin)
<b>Facet</b> <sup>®</sup> 75 DF (quinclorac)	<b>Sinbar</b> <sup>®</sup> (terbacil)
<b>FirstRate</b> <sup>®</sup> (cloransulam-methyl)	<b>Stinger</b> <sup>®</sup> (clopyralid)
<b>Flexstar</b> <sup>®</sup> (fomesafen)	<b>Storm</b> <sup>®</sup> (bentazon + acifluorfen)
<b>Glyphosate</b>	<b>Synchrony</b> <sup>®</sup> XP (chlorimuron + thifensulfuron)
<b>Londax</b> <sup>®</sup> (bensulfuron)	<b>Thifensulfuron</b>
<b>Liberty</b> <sup>®</sup> (glufosinate)	<b>Thistrol</b> <sup>®</sup> (MCPB)
<b>Lightning</b> <sup>®</sup> (imazethapyr + imazapyr)	<b>Ultra Blazer</b> <sup>®</sup> (acifluorfen)
<b>MCPA</b>	<b>2,4-DB</b>
<b>Outlook</b> <sup>®</sup> (dimethenamid-P)	

Tank mixes must be combined and applied in accordance with the directions in this label and the labels of all tank mix partners. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If all target weeds species are not at the specified stage of growth for the timing of treatment at the same time, make separate applications.

Mixing this product with other registered fertilizer, additives or pesticide products (insecticides, fungicides, miticides or herbicides) may result in physical incompatibility of the products, crop injury or a reduction in weed control. Applicators should only tank mix this product with those products listed in this label. When mixing this product with products not listed in this label, consult local agricultural authorities for information.

### Compatibility Test for Mix Components

Before mixing additives and/or other pesticides, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly.

Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of specified label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

## Mixing Order

When mixing additives and/or other pesticides in a spray tank, add the products to be used in the following sequence:

1. Water - Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2. Agitation - Maintain constant agitation throughout mixing and application.
3. Products in PVA Bags - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. Water-dispersible Products - Such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions. If an inductor is used, rinse it thoroughly after the component has been added.
5. Water-soluble Products - Such as **Biscayne**. If an inductor is used, rinse it thoroughly after the component has been added.
6. Emulsifiable Concentrates - Such as oil concentrate when applicable. If an inductor is used, rinse it thoroughly after the component has been added.
7. Water-soluble Additives - Such as AMS or UAN when applicable. If an inductor is used, rinse it thoroughly after the component has been added.
8. Remaining quantity of water.

NOTE: Maintain constant agitation during application.

## RESTRICTIONS AND LIMITATIONS THAT APPLY TO ALL CROPS

- Do NOT exceed the maximum seasonal use rate of 4 pints of **Biscayne** per acre (2 pounds active ingredient), per season, from all sources.
- Do NOT apply more than 3 pints of **Biscayne** per acre in a single application (except for treatment of kochia and hairy nightshade on Peppermint and Spearmint. Please see specific crop restrictions).
- The minimum permitted retreatment interval for bentazon products is 7 days (except for treatment of kochia and hairy nightshade on Peppermint and Spearmint. Please see specific crop restrictions).
- **Biscayne** must NOT be applied through an irrigation system of any type.
- Do NOT enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 48 hours.
- Do NOT apply **Biscayne** if crops in the treatment area are suffering stress due to flooding, drought, widely varying temperatures, hail damage or injury from previously applied pesticide(s) as crop damage may result.
- Do NOT apply **Biscayne** to crops injured by other pesticide product treatment(s) (e.g. plant stunting or phytotoxicity) as previously caused injury may be prolonged and/or enhanced.

## PRECAUTIONS THAT APPLY TO ALL CROPS

- Unsatisfactory control of target species may result if **Biscayne** is applied to weeds suffering stress caused by injury from previously applied pesticide(s), drought, other mechanical injury or cold temperatures. Overhead irrigation or precipitation within 4 hours of treatment with this product may reduce the effect of **Biscayne** in controlling target weed species.



## CROP SPECIFIC DIRECTIONS

Apply **Biscayne** during early postemergence. Treatment with this product must take place before target weeds reach the maximum growth stage for application listed in the **Application Rates for Specific Weed Growth Stages for All Crops EXCEPT Rice** table above. For application rates for rice crops, see the **RICE** section below.

### BEANS

#### (DRY AND SUCCULENT)

The following beans are tolerant of **Biscayne**:

- Adzuki Beans
- Black Turtle Soup Beans
- Cranberry Beans
- Great Northern Beans
- Kidney Beans
- Lima Beans (Small & Large)
- Navy Beans
- Pink Beans
- Pinto Beans
- Red Beans
- Snap Beans
- White Beans

Beans crops are tolerant to applications of **Biscayne** once the first trifoliolate leaf has fully expanded. However, even when the bean crop is in the tolerant stage of growth there may be bronzing, yellowing, burning, speckling or burning of leaves under certain conditions (see **Restrictions** below). Such crop injury is temporary and the plant will outgrow it without affecting podset, maturity and without reducing yield. Use of oil with **Biscayne** may have the effect of reducing yield and increasing crop injury.

#### Restrictions

- Do not apply **Biscayne** on its own to succulent or dry beans grown in South Carolina or Georgia.
- When applying **Biscayne** to succulent or dry beans grown in South Carolina or Georgia, this product must be combined with Pursuit herbicide or Raptor herbicide. Apply **Biscayne** at a rate of 6 - 16 fl. oz. per acre. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not treat field beans with **Biscayne** until they have a minimum of one trifoliolate leaf fully expanded.
- In order to avoid severe crop damage, **Biscayne** must not be applied to:
  - blackeyes in California
  - garbanzo beans at any growth stage
  - lupines at any growth stage
- **Biscayne** must not be applied to succulent or dry beans within 30 days of harvest.
- For control of yellow nutsedge in California, treat with 2 pints of **Biscayne** per acre when plants are 6 - 8 inches tall. 10 to 14 days after the first application, make a second treatment at the same rate.

#### Precautions

- When applying **Biscayne** to snap beans, the use of an oil additive may enhance the potential for leaf burn and crop injury.
- **Biscayne** is not recommended for use on adzuki beans in California.

#### Tank Mixes

**For Dry Beans**, **Biscayne** may be tank mixed with one of the following herbicides: Outlook<sup>®</sup>, Poast<sup>®</sup>, Pursuit<sup>®</sup> or Raptor<sup>®</sup>.

**For Succulent Beans**, **Biscayne** may be tank mixed with one of the following herbicides: Poast<sup>®</sup> or Pursuit<sup>®</sup>. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## CLOVER GROWN FOR SEED

### (OREGON AND WASHINGTON)

Apply **Biscayne** as a postemergence treatment for clover grown for seed in Oregon and Washington. Make a foliar broadcast application in the spring at up to 2 pints of this product per acre. A second application 7 to 14 days later can be made if required at the same use rate.

Clover is tolerant to treatment with **Biscayne**. Under certain conditions, leaf burn may take place, however, within 10 days, clover will generally outgrow the condition.

Add a crop oil concentrate (COC) that is nonphytotoxic as directed in the **Additive Rates** table in the **ADDITIVES** section at the beginning of this label.

#### Restrictions

- Do not allow treated areas to be used to harvest forage, hay or feed for livestock feed or for livestock grazing for at least 36 days after an application of **Biscayne**.

## CORN

### (INCLUDING CORN GROWN FOR SEED/SILAGE, FIELD, POPCORN, SWEET) AND SORGHUM (INCLUDING FORAGE AND GRAIN)

Producers of seeds must refer to the seed company for information on seed production inbred lines' tolerance to applications of **Biscayne**.

#### Restrictions

- For sorghum, do not exceed 2 pints of **Biscayne** per acre per year.
- Do not apply **Biscayne** to sorghum that is blooming or heading.
- Do not allow grazing in treated areas for 12 days after the treatment with this product.
- **Biscayne** must not be used on forage sorghum in California.

#### Precautions

- **Biscayne** is not recommended for treatment of sorghum or corn in California to control yellow nutsedge.

#### Tank Mixes

Do NOT tank mix this product with atrazine in California.

**For Corn Applications**, **Biscayne** may be tank mixed with one of the following products (including herbicides that are registered for use in corn hybrids that are tolerant to treatment with glufosinate, glyphosate and imidazolinone): Atrazine, Atrazine + Dicamba, Clarity<sup>®</sup>, Distinct<sup>®</sup>, Glyphosate, Outlook<sup>®</sup>, Liberty<sup>®</sup>, Lightning<sup>®</sup>, or Pursuit<sup>®</sup>.

**For Sorghum Applications**, **Biscayne** may be tank mixed with one of the following products: Atrazine, Atrazine + Dicamba, Clarity<sup>®</sup>, Outlook<sup>®</sup>, or Paramount<sup>®</sup>.

## PEPPERMINT AND SPEARMINT

Although, spearmint and peppermint are tolerant to applications with this product, treatment may cause leaf-burning. This may occur when crops are actively growing and have new, succulent tissue. Crops will generally grow out of the condition within 10 days.

In order to control kochia and hairy nightshade, **Biscayne** may be applied at a rate of 4.0 pints per acre in one application. In order to control kochia, combine **Biscayne** with an oil concentrate.

#### Tank Mixes

**Biscayne** may be tank mixed with one of the following products: Buctril<sup>®</sup>, Poast<sup>®</sup>, Sinbar<sup>®</sup> or Stinger<sup>®</sup>. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## PEAS

### (DRY AND SUCCULENT)

Peas (English, garden and Southern) are tolerant of applications of **Biscayne** once 4 nodes or 3 pairs of leaves are present. Crop injury may occur under certain conditions such as bronzing, yellowing, burning or speckling. Such damage will be temporary and the crop will generally outgrow it without causing any reduction in yield or delaying podset/maturity.

Do not apply **Biscayne** in Western irrigated areas during 2 - 5 day periods of cold weather (i.e. temperature below 75°F during the day below 55°F during the night). Applications during a prolonged cold spell may nullify weed control.

#### Restrictions

- Do not apply **Biscayne** on its own in South Carolina and Georgia to succulent peas.
- When treating succulent peas grown in South Carolina and Georgia, **Biscayne** must be tank mixed with Pursuit herbicide or Raptor herbicide. Apply **Biscayne** at a rate of 6 - 16 fl. oz. per acre. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not apply **Biscayne** to dry peas within 30 days of harvest.
- When treating succulent peas, do not apply **Biscayne** within 10 days of harvest.
- When treating succulent peas in California, do not apply **Biscayne** within 30 days of harvest.
- Do not treat peas that are suffering stress from root rot.
- Do not treat pea crops when they are in bloom.
- Do not treat the following crops with **Biscayne** at any stage of growth:
  - blackeyes grown in California
  - garbanzo beans
  - lupines
- Oil must not be added to **Biscayne** except when treating peas in the Pacific Northwest.

#### Precautions

- There may be a higher likelihood of crop damage from applications with **Biscayne** if there are in-furrow treatments of nematocide or insecticide.

#### Tank Mixes

NOTE: The following tank mixes are not applicable in California. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Biscayne** may be tank mixed with one of the following products: \*MCPA, Pursuit®, Raptor® or \*\*Thistrol®.

\* In order to enhance control of common lambsquarters and pigweed species, applicators may use a tank mix of **Biscayne** plus MCPA.

\*\* Tank mixing with Thistrol® is for use in ME, NH, VT, MA, CT, RI, NY, PA, NJ, VA, MD, DE, WA, ID, and OR. Apply after the 4 node/3-leaf stage and no later than 3 nodes before peas flower. Because of the variation among pea cultivars and treatment, producers and sellers have not determined the safety of a combination of **Biscayne** and Thistrol® for use in all conditions on all pea crops. Applicators must therefore determine whether this tank mix can be used safely before any broader treatment.

#### Tank Mix Restrictions

- Do NOT use any oil-based additives or any other surfactants or spray additives with the above tank mixes.
- Do NOT treat peas with the above tank mixes when temperatures are greater than 90°F.
- Do NOT treat peas with the above tank mixes once pea flower buds have emerged.

#### Tank Mix Precautions

- The following crops are particularly sensitive to Thistrol:
  - Beans
  - Cotton
  - Grapes
  - Ornamentals
  - Tomatoes

## PEANUTS

Apply **Biscayne** to peanuts from peanut cracking through pegging.

In-furrow applications of nematocides and/or insecticides may increase the likelihood of crop injury from treatment with **Biscayne**.

Hay and forage from peanut crops may be used as livestock feed.

#### Restrictions

- Do NOT allow grazing in treated fields for a minimum of 50 days after **Biscayne** is last applied.

#### Tank Mixes

NOTE: The following tank mixes are not applicable in California. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Biscayne** may be tank mixed with one of the following products: Avalon, Ultra Blazer®, Outlook®, Poast®, \*Paraquat, or 2,4-DB amine.

\* Apply this tank mix combination at the ground crack stage of growth in order to control early weed flushes. A second treatment may be applied up to 28 days after ground crack stage. Applicators must mix this tank mixture with a nonionic surfactant containing at least 50% surface active agent at specified rates.

#### Tank Mix Restrictions

- Do NOT combine ammonium sulfate or a UAN solution with a **Biscayne**/Avalon, Ultra Blazer/Poast tank mix.
- Do NOT use with a tank mix of **Biscayne** plus Paraquat.
- Do NOT use any oil based additive or UAN with a tank mix of **Biscayne** plus 2,4-DB.
- Only use amine formulations of 2,4-DB.

## RICE

### (Not for use in California)

Treat with **Biscayne** as an early postemergence application, prior to target weeds reaching the growth stages listed in the **Application Rates for Rice** tables below. Straw from rice may be used to feed livestock.

#### Application Instructions

In order to achieve the best coverage with **Biscayne**, orient nozzles straight back. Do not place nozzles further out than three quarters of the distance from the center of the aircraft to the end of the rotor or wing.

#### Alternate Flooding Culture

In TX, LA, AR, and MS, the weed growth of target species corresponds generally to tillering (stooling) rice before the field is permanently flooded.

Apply **Biscayne** a minimum of 24 hours prior to flooding when there is no water on the treatment area.

If application of **Biscayne** is not possible prior to flooding, refer to the **Continuous Flooding Culture** section below.

## Continuous Flooding Culture

In states where continuous flooding culture is used, or when **Biscayne** is applied after permanent flooding, apply this product when target species are above the water surface. Control of target species that are under water when this product is applied will be inadequate.

In order to make an early application of **Biscayne**, water may be completely or partially drained in order to expose target species to treatment. The water level must not be raised for a minimum of 24 hours after treatment. Early flooding following application may result in inadequate control.

Applicators must not use ground equipment to treat flooded fields. The resultant splashing will wash **Biscayne** off the surface of the weed resulting in inadequate control.

## Restrictions

- Do not apply **Biscayne** in a field where there is commercial cultivation of crayfish or catfish.
- Water treated with this product or containing residue of **Biscayne** from rice must not be used for the irrigation of crops that are not labeled for treatment with **Biscayne**.
- Do not exceed 4 pints of **Biscayne** per acre per season. This applies whether there are one or two rice crops grown that season (including ratoon).

## Application Rates for Rice – Flooded Fields

Weeds Controlled	Application Rate (Pints)	Max. Height Above Soil (Inches)	Height Range Above Water Level (Inches)	Comment
Cocklebur	1.5	10	3 - 6	If after the first application of <b>Biscayne</b> a second weed flush occurs, re-apply this product to the treatment area in accordance with this table.
	2.0	15	6 - 10	
Dayflower	1.5	6	3 - 5	
	2.0	10	5 - 8	
Redstem	1.5	4	2 - 3	
	2.0	8	4 - 6	
Smartweed	1.5	6	2 - 5	
	2.0	10	5 - 8	
Water Plantains, Arrowhead	1.5	Not Rec.	-	
	2.0	7	5 - 6	
Water Plantains, Common	1.5	Not Rec.	-	
	2.0	7	5 - 6	
Yellow Nutsedge	1.5	6	4 - 5	
	2.0	10	6 - 8	

## Application Rates for Rice – Drained Fields

Weeds Controlled	Application Rate (Pints)	Leaf Stage (Leaves)	Max. Height (Inches)	Comment
Cocklebur	1.5	2 - 10	10	If after the first application of <b>Biscayne</b> a second weed flush occurs, re-apply this product to the treatment area in accordance with this table.
	2.0	10 - 15	15	
Dayflower	1.5	2 - 10	6	
	2.0	10 - 15	10	
Ducksalad	1.5	Not Rec.	-	
	2.0	6 - 10	6	
Eclipta	1.5	4 - 6	2	
	2.0	4 - 6	2	
Gooseweed	1.5	4 - 6	4	
	2.0	6 - 10	8	
Redstem	1.5	up to 6	4	
	2.0	6 - 10	8	
Redweed	1.5	4 - 6	6	
	2.0	6 - 10	8	
Smartweed	1.5	2 - 10	6	
	2.0	10 - 15	10	
Spikerush	1.5	2 - 6	6	
	2.0	6 - 8	8	
Water Plantains, Arrowhead	1.5	Not Rec.	-	
	2.0	up to 4	7	
Water Plantains, Common	1.5	Not Rec.	-	
	2.0	up to 4	7	
Yellow Nutsedge	1.5	4 - 6	6	
	2.0	6 - 8	10	

## Tank Mixes

**Biscayne** may be tank mixed with one of the following products: Avalon, Ultra Blazer<sup>®</sup>, Facet<sup>®</sup> 75 DF, Londax<sup>®</sup>, Propanil, \*Storm<sup>®</sup>.

\*Apply this tank mix with the labeled rate of Storm combined with 0.5 - 1.0 pint of **Biscayne** per acre after the 3-leaf stage. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## Tank Mix Restrictions

- The **Biscayne** plus Londax<sup>®</sup> herbicide tank mixture must be applied within 7 days of the establishment of permanent flood.
- The **Biscayne** plus propanil tank mix must only be applied to drained fields.
- Do not use crop oil concentrate in tank mixes with propanil. Tank mix this product with propanil according to the active ingredient content in the product used. Propanil tank mixes must be tested for physical compatibility with **Biscayne** prior to application.

## SOYBEANS

Although soybeans are tolerant at all stages of growth to applications of **Biscayne**, under certain conditions, slight leaf bronzing and speckling may occur. Soybean crops will outgrow these conditions in general, within 10 days.

### Restrictions

- Do not allow grazing on treated areas for 30 days (minimum) after the last application of **Biscayne**.
- Do not cut treated soybean for forage or hay for 30 days (minimum) after the last application of **Biscayne**.

### Tank Mixes

NOTE: The following tank mixes do not apply to California.

**Biscayne** may be tank mixed with one of the following products\*: Avalon, \*\* Chlorimuron + Thifensulfuron, Classic®, Cobra®, \*\*Concert® II, \*\*FirstRate®, Flexstar®, Outlook®, Liberty®, Poast®, Poast Plus®, Pursuit®, Raptor®, Reflex®, Resource®, Roundup® Ultra, Scepter®, Synchrony® XP, \*\*Thifensulfuron, Ultra Blazer® or 2,4-DB amine.

\* Includes Roundup Ready®, LibertyLink® and STS™ varieties. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

\*\*UAN at a rate of 2 - 4 pints per acre and a nonionic surfactant at a rate of 1 - 2 pints per 100 gallon are recommended for these tank mixtures.

Treatment with this tank mix will result in crop damage (e.g. bronzing, crinkling or burning) which may cause a reduction in yield.

### Tank Mix Restrictions

- **Biscayne** plus Avalon or Ultra Blazer plus Poast: an oil concentrate must be used with this tank mix instead of a spray surfactant.
- **Biscayne** plus Chlorimuron + Thifensulfuron: Do not use an oil concentrate with this tank mix except in soybean varieties designated as STS.

### Restrictions: Biscayne plus 2,4-DB Amine Tank Mix

- Only use 2,4-DB that is an amine formulation, and do not use any adjuvant, other than UAN at a rate of 2 - 4 pints per acre with this tank mix.
- Do not exceed 1 treatment of this tank mix per season.
- Do not treat soybeans that show signs of disease such (e.g. phytophthora root rot) with this tank mix.

### Mixing Biscayne with Insecticides

If foliar control or postemergence control of certain insects is required in the soybean crop, **Biscayne** may be combined with an insecticide. In order to do so, the specified application time of the insecticide product must coincide with the specified application time of this product.

Restriction: **Biscayne** must NOT be tank mixed with Sevin® or malathion insecticides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Biscayne** may be tank mixed with one of the following insecticides: dimethoate, Pounce®, Pydrin® or Lorsban® 4E.

Applying a tank mix of this product plus an insecticide may increase the likelihood of crop damage. The conditions in which this product is mixed with an insecticide product may vary. Certain conditions may reduce mixing quality. Test the proposed combination of insecticide and **Biscayne** in accordance with the **Compatibility Test for Mix Components** section, prior to application.

## COMMERCIAL TURF & ORNAMENTAL AND NONCROPLAND USES

### USE INSTRUCTIONS

This product is a selective herbicide for control of annual sedges, broadleaf weeds, and yellow nutsedge. Apply **Biscayne** as a postemergence treatment in the following use sites: ornamentals, established turfgrass, noncropland sites, nurseries, rights-of-way and roadsides.

Read and follow the specific use site directions below for each site use.

**Biscayne** works mainly through contact with target species. Therefore, ensure all weeds are covered thoroughly with the spray application.

Overhead sprinkler irrigation or rainfall may reduce the effectiveness of an application of **Biscayne** if it occurs within 8 hours of treatment.

### Application Instructions

Clean application equipment thoroughly prior to application to prevent potential crop injury from previously applied products.

Apply **Biscayne** to actively growing weeds as a postemergence treatment. Target species that are growing under conditions of drought may be inadequately controlled. If moisture content in the soil is insufficient for weeds to be growing actively, irrigate prior to application. **Biscayne** does not provide control of grass weeds.

Apply spray with a hose-end type sprayer, a handheld pump-up or a knap-sack sprayer with standard high-pressure flat fan or hollow-cone nozzles that are spaced 20-inches apart. Do not apply this product with whirl chamber, controlled droplet applicator (CDA), or flood nozzles.

Apply **Biscayne** in 1 gallon of water (minimum) per 1000 sq. ft. (40 gallons per acre). Use a spray pressure of at least 40 psi. Spray pressure must be measured at the boom, not in the line or at the pump.

Where there is dense foliage or dense weed coverage, use a higher water volume (2.5 gallons per 1000 sq. ft.) and increased spray pressure (80 psi).

### Mixing Instructions

Use new spray mixture for each treatment, and mix sufficient solution for just one application.

Fill clean spray application equipment tank 1/2 - 2/3 full with clean water and start agitation prior to adding the specified amount of **Biscayne**. Ensure the spray solution mixes thoroughly. Next, add oil concentrate and the rest of the water. Maintain agitation throughout mixing and treatment.

Mixing **Biscayne** with other additives, fertilizers or pesticides (including insecticides, herbicides, fungicides and miticides) may cause physical incompatibility between tank mix partners, a reduction in weed control, or damage to ornamentals and/or turfgrass. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Use Rates

- Do NOT exceed 0.75 fl. oz. of **Biscayne** per 1000 sq. ft. in a single application.
- Do NOT exceed 32 fl. oz. (1 quart) of **Biscayne** per acre in a single application (1.0 lb. of active ingredient).
- Do NOT exceed 64 fl. oz. (2 quarts) of **Biscayne** per acre in a season (2.0 lb. of active ingredient).

Apply **Biscayne** with an oil concentrate (see next section) at the following rates (see the **Application Rates for Commercial Turf and Ornamentals** table below for specific rates based on size and type of target weeds):

For treatment areas of 1000 sq. ft., apply at a rate of 0.55 - 0.75 fluid ounces in 1 - 2 gallons of water.

For treatment areas of 1 acre, apply at a rate of 12 - 16 fluid ounces in 40 - 80 gallons of water.

**IMPORTANT:** Use of the above rates in a spot spray treatment for individual weeds may lead to excessive dosage and possible damage to ornamentals or turfgrass.

### Oil Concentrate

For optimal control of target species, an oil concentrate must be added to the spray solution. The oil concentrate must contain a vegetable-oil or a petroleum-oil base. Suitable examples are crop oil concentrate (COC) and methylated seed oil (MSO). The selected oil concentrate must be nonphytotoxic, must contain only ingredients that are EPA exempt and must have been successful according to local experience. Vegetable-oil and petroleum-oil based concentrates should contain emulsifiers in order to provide good mix properties. Highly refined vegetable oil based concentrates are likely to mix better than those that are unrefined.

Apply oil concentrate in the application spray solution at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).

Applying **Biscayne** with an oil concentrate under certain conditions (e.g. when temperature and relative humidity are high) may cause slight leaf burn to turfgrass, ornamentals and other desired plants. Talk to your **Biscayne** supplier in order to assess the local success of potential oil concentrates.

Do not add oil concentrate to a tank mix that includes a product/products whose label(s) prohibit or caution against their application with oil adjuvants.

In addition, the oil concentrate selected must mix adequately with the spray solution in the following jar test:

### Jar Test to Estimate Oil Concentrate Suitability

Use the following jar test to determine the suitability of an oil concentrate:

1) Add the ingredients of the tank mix in the following order and gently mix in between adding ingredients:

- Water
- **Biscayne**
- Tank mix partner (if used)
- Oil concentrate

Use water from the intended source for the application solution at source temperature.

If the intended application volume is 1 gallon per 1000 sq. ft., use 1600 ml of water. Adjust the test water volume proportionately if there is a different intended application water volume.

2) Add 2 teaspoons of oil concentrate and 2 teaspoons of **Biscayne** for every 0.75 fl. oz. per 1000 sq. ft. in the intended application solution at the specified rate.

3) Once all the ingredients have been added, put a lid on the jar. Invert the jar 10 times to mix the solution, then leave the solution to sit for 15 minutes.

If the tank mix displays any of the following features, it may not be suitable for application:

- Globules or a film of oil on the surface of the solution.
- Tank mix solution is not uniform.
- Fine particles suspended in the liquid (known as flocculation). Particles may also appear at the bottom of the jar as a precipitated layer.
- If the texture of the solution thickens (clabbering) so that it is coagulated with a similar consistency to yogurt or cottage cheese.

### Restrictions

- Do not exceed 32 fl. oz. (1 quart) of **Biscayne** per acre in a single application (1.0 lb. of active ingredient).
- Do not exceed 64 fl. oz. (2 quarts) of **Biscayne** per acre in a season (2.0 lb. of active ingredient).
- Do not apply this product in conditions that favor drift. Contact with adjacent plants or crops may cause damage.
- Do not apply **Biscayne** to open waters.

### Weeds Controlled

Treat actively growing target weed species with **Biscayne** as an early postemergence application. Weeds must be treated before they reach the maximum growth stage/height listed in the **Application Rates for Commercial Turf and Ornamentals** table below. For optimal treatment, apply **Biscayne** to small or newly emerged weeds. Later treatment may allow target species to grow beyond the specified growth stage/height which will result in reduced or inadequate control (except musk thistle, Canada thistle and yellow nutsedge - refer to **special instructions** below).

### Application Rates for Commercial Turf and Ornamentals

Weeds Controlled	Application Rate per 1000 sq. ft. (fl. oz.)	Leaf Stage (Leaves)	Max. Height (Inches)	Comment
Anoda, spurred <i>Anoda cristata</i>	0.55 0.75	up to 6 6 - 8	3 4	
Balloonvine <i>Cardiospermum halicacabum</i>	0.55 0.75	2 - 4 4 - 6	2 3	
Buckwheat, wild <i>Polygonum convolvulus</i>	0.55 0.75	up to 4 4 - 6	3 5	
Coffee senna <i>Cassia occidentalis</i>	0.55 0.75	Not Rec. up to 1 pinnate*	- 2	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Dayflower <i>Commelina</i> spp.	0.55 0.75	up to 6 6 - 10	4 8	
Devil's-claw <i>Proboscidea louisiana</i>	0.55 0.75	Not Rec. up to 6*	- 2	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Galinsoga <i>Galinsoga</i> spp.	0.55 0.75	Not Rec. Cotyledon to 6*	- 2	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Groundsel, Common <i>Senecio vulgaris</i>	0.55 0.75	Not Rec. 2 - 10	- 6	

(continued)

**Application Rates for Commercial Turf and Ornamentals (continued)**

Weeds Controlled	Application Rate per 1000 sq. ft. (fl. oz.)	Leaf Stage (Leaves)	Max. Height (Inches)	Comment
Ladysthumb <i>Polygonum persicaria</i>	0.55 0.75	up to 6 6 - 10	6 10	
Lambsquarters, common <i>Chenopodium album</i>	0.55 0.75	Not Rec. 4 - 8*	- 2	Control of this species may be inconsistent/partial. *Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Mustard, wild <i>Sinapis arvensis</i>	0.55 0.75	up to 6 6 - 10	4 8	
Nutsedge, yellow <i>Cyperus esculentus</i>	0.55 0.75	See <b>Comment</b>		Yellow nutsedge emerges all through the year in southern USA. It emerges from May through July in northern USA. <b>Biscayne</b> can only control emerged weeds, therefore make the first applications of this product when the weed emerges. For the most effective control of yellow nutsedge, make 2 treatments with <b>Biscayne</b> at a rate of 0.55 - 0.75 fl. oz. per 1000 sq. ft. (12 - 16 fl. oz. per acre). Apply this product when the weed is 6 - 8 inches tall. Treat a second time when new growth appears later in the season or 7 - 10 days after the first application. Use the same use rate as the first application. In order to achieve optimal control of yellow nutsedge, thorough spray coverage is essential.
Poinsettia, wild <i>Euphorbia heterophylla</i>	0.55 0.75	2 - 4 4 - 8*	4 6	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).

(continued)

**Application Rates for Commercial Turf and Ornamentals (continued)**

Weeds Controlled	Application Rate per 1000 sq. ft. (fl. oz.)	Leaf Stage (Leaves)	Max. Height (Inches)	Comment
Prickly sida/ Teaweed <i>Sida spinosa</i>	0.55 0.75	up to 6 6 - 8	3 4	
Purslane, common <i>Portulaca oleracea</i>	0.55 0.75	up to 4 4 - 6	1 2	
Ragweed, common <i>Ambrosia artemisiifolia</i>	0.55 0.75	Not Rec. 4 - 6*	- 3	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Ragweed, giant <i>Ambrosia trifida</i>	0.55 0.75	Not Rec. up to 4	- 6	If there is a second flush of weeds following the first treatment, make a second application in accordance with the rates in this table.
Redweed <i>Melochia corchorifolia</i>	0.55 0.75	4 - 6 6 - 10	6 8	
Sedge, annual <i>Cyperus compressus</i>	0.55 0.75	Not Rec. Not Rec.	6 - 8 6 - 8	
Sesbania <i>Sesbania exaltata</i>	0.55 0.75	Not Rec. 3 - 5*	- 3	*Requires use of an oil concentrate at a rate of 0.75 fl. oz. per 1000 sq. ft. (2 pints per acre).
Shepherd's purse <i>Capsella bursa-pastoris</i>	0.55 0.75	up to 6 6 - 10	4 8	Do not apply to the rosette prior to seed stalk emerging.
Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>	0.55 0.75	up to 6 6 - 10	6 10	
Spurweed/Lawn burrweed <i>Soliva pterosperma</i>	0.55 0.75	Not Rec. 2 - 6	- 3	
Sunflower, wild <i>Helianthus annuus</i>	0.55 0.75	up to 4 4 - 6	5 8	

(continued)

## Application Rates for Commercial Turf and Ornamentals (continued)

Weeds Controlled	Application Rate per 1000 sq. ft. (fl. oz.)	Leaf Stage (Leaves)	Max. Height (Inches)	Comment
Thistle, Canada <i>Cirsium arvense</i>	0.55 0.75	* *	* *	Apply <b>Biscayne</b> to Canada thistle when the weeds are between 8 inches tall and the bud stage of growth at a rate of 0.75 fl. oz. per 1000 sq. ft. (16 fl. oz. per acre). If adequate control is not achieved with the first application, apply this product again when new growth appears, or 7 - 10 days after the first application at the same rate.
Thistle, musk <i>Carduus nutans</i>	0.55 0.75	* *	* *	Apply <b>Biscayne</b> to musk thistle at a rate of 0.75 fl. oz. per 1000 sq. ft. (16 fl. oz. per acre). Treat with this product when the weed is in the rosette stage and has a diameter no greater than 10 inches. If adequate control is not achieved by the first treatment, apply this product again when new growth appears or 7 - 10 days after the first application. Use the same use rate as the first application.

### INSTRUCTIONS FOR ESTABLISHED TURFGRASS

Treat established turfgrass growing in areas such as sod farms, commercial or residential settings, recreational areas, golf courses, athletic fields or any other area of established or maintained turfgrass. **Biscayne** may be used to treat the following established turfgrass species:

- Bahiagrass
- Bentgrass
- Bermudagrass
- Bluegrass
- Buffalograss
- Carpetgrass
- Centipedegrass
- Fescue
- Ryegrass
- St. Augustinegrass
- Zoysiagrass

Refer to the **Application Rates for Commercial Turf and Ornamentals** table above for a complete list of weeds controlled.

### Restrictions

- Do not treat turfgrass that has suffered stress, for example from cold temperature, injury from other pesticides or drought.
- **Biscayne** must not be applied to newly sprigged or newly seeded turfgrass until the sprigs or seedlings are well established.
- Do not apply **Biscayne** to golf course greens or collars.
- Do not exceed 0.75 fl. oz. of **Biscayne** per 1000 sq. ft. (32 fl. oz. per acre) in perennial ryegrass in a single application. Do not make further applications for at least 21 days.
- Make the first treatment with **Biscayne** in unmowed established turfgrass after emergence but when yellow nutsedge, Canada thistle and annual sedge are less than 8-inches in height. Treat annual broadleaf weeds when they are 4 inches in height or less.
- Do not mow treated turfgrass within 3 days after or before treatment with this product in order to ensure an optimal application to control broadleaf weeds.
- Do not mow turfgrass for 5 days (minimum) after treatment for optimal control of sedges.

### Precautions

- When treating turfgrass with this product, avoid over-the-top spraying of adjacent ornamental trees, shrubs, and flowers unless otherwise specified in this label. Application of **Biscayne** at the base of established ornamental shrubs, trees and flowers should not cause damage except for in rhododendron and sycamore.

### Tank Mixes for Established Turfgrass

For postemergence control of other sedges and broadleaf weeds that are not specified in this label, tank mix **Biscayne** with other registered products labeled for use in turfgrass. For example:

- Turflon® herbicide
- Image® 70 DG herbicide
- MSMA
- 2,4-D
- Atrazine
- mixes of 2,4-DP (dichlorprop) or 2,4-D, MCPP (mecoprop)

**IMPORTANT:** Some of these products cannot be used on all turfgrass sites/species. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank mix **Biscayne** with Segment® herbicide for treatment of fine fescue and centipede grass species.

Applicators must determine the compatibility of tank mix partners prior to application. Applicators can use an anti-foaming agent if required.

Applicators must NOT use an oil additive or a surfactant with 2,4-DP; 2,4-D; or MCPP.

When making tank mixes other than those specified in this label, refer to local professional authorities. Also, test the tank mix in a small area and allow 7 - 10 days to ascertain potential damage prior to wider application.

## INSTRUCTIONS FOR ORNAMENTALS, NURSERY, NONCROPLAND SITES, ROADSIDES, AND RIGHTS-OF-WAY

Treat the following ornamental species with **Biscayne** as an over the top application:

Alumroot ( <i>Heuchera</i> spp.)	Holly, dwarf Chinese ( <i>Ilex cornuta</i> )
Apple (nonbearing) ( <i>Malus</i> spp.)	Holly, Japanese ( <i>Ilex crenata</i> )
Arborvitae <sup>1</sup> ( <i>Thuja occidentalis</i> )*	Hydrangea ( <i>Hydrangea</i> spp.)
Barberry, Japanese ( <i>Berberis thunbergii</i> )	Jasmine ( <i>Jasminum</i> spp.)
Boxwood ( <i>Buxus</i> spp.)	Lily, plantain ( <i>Hosta fortunei</i> )
Bugle, common ( <i>Ajuga</i> spp.)*	Lilyturf ( <i>Liriope</i> spp.)
Butterfly bush ( <i>Buddleia davidii</i> )	Lilyturf, big blue ( <i>Liriope muscari</i> )
Cabbage ornamental ( <i>Brassica</i> spp.)	Liriope, creeping ( <i>Liriope spicata</i> )
Cape jasmine ( <i>Gardenia</i> spp.)	Marigold ( <i>Tagetes</i> spp.)
Chokeberry ( <i>Photinia</i> spp.)*	Myrtle, wax ( <i>Myrica cerifera</i> )
Coral bells ( <i>Heuchera</i> spp.)*	Oak, red <sup>1</sup> ( <i>Quercus rubra</i> )*
Cotoneaster ( <i>Cotoneaster</i> spp.)	Pachysandra ( <i>Pachysandra terminalis</i> )*
Crabapple (nonbearing) ( <i>Malus</i> spp.)*	Petunia ( <i>Petunia hybrid</i> )
Crape myrtle ( <i>Lagerstroemia indica</i> )	Photinia ( <i>Photinia</i> spp.)*
Cypress, bald ( <i>Taxodium distichum</i> )	Pine, Mugo ( <i>Pinus mugo</i> )
Daylily ( <i>Hemerocallis</i> spp.)	Pine, white ( <i>Pinus strobus</i> )
Dusty miller ( <i>Centaurea cineraria</i> )	Pittosporum, Japanese ( <i>Pittosporum tobira</i> )
Euonymus ( <i>Euonymus</i> spp.)	Snapdragon ( <i>Antirrhinum majus</i> )
Gardenia, common ( <i>Gardenia</i> spp.)	Yew ( <i>Taxus</i> spp.)
Golden-rain tree ( <i>Koelreuteria bipinnata</i> )	Yew hybrids ( <i>Taxus x media</i> )*
Hawthorn, Indian ( <i>Raphiolepis indica</i> )	Yew, Japanese ( <i>Taxus cuspidata</i> )*
Holly ( <i>Ilex</i> spp.)	Yew, Southern ( <i>Podocarpus macrophyllus</i> )
Holly, Chinese ( <i>Ilex cornuta</i> )	

<sup>1</sup> Do not exceed one application per growing season per crop.

\* Not for use in California.

Some species may display different levels of tolerance to an application of **Biscayne**. Make a test application of this product on a small number of plants of the species to which a wider application will be made, before wider application. Assess the effect of the application over 2 weeks before wider application.

Due to variation in application techniques and within species, the safety of this product for use on all nursery plants/ornamentals has not been completely determined under all conditions of growth. Applicators must therefore test a small area/a few plants in order to assess suitability of use prior to wider application.

When treating all other ornamental and landscape shrubs, trees, nursery plants and flowers not listed above, make a directed spray application with **Biscayne** directed away from the foliage of desired plants.

Applications of this product may be made where grass vegetation needs to be maintained.

### Restrictions

- Do NOT use an oil concentrate with **Biscayne** when making an over the top application in ornamentals.
- Do not apply this product to ornamental or nursery plants that have been subject to stress conditions such as hail damage, flooding, drought, extreme heat, or widely fluctuating temperatures.
- Do not apply **Biscayne** if ornamental or nursery plants show injury (leaf phytotoxicity or plant stunting) produced by prior herbicide applications.
- Do not treat with this product as a directed spray over the roots or under the tree line of rhododendron or sycamore, or damage may result. Do not treat with this product if the risk of damage to rhododendron or sycamore is unacceptable.
- When making application to Arborvitae (*Thuja occidentalis*) or Red Oak (*Quercus rubra*), do not exceed one application per growing season.

### Tank Mixes

#### Biscayne plus Segment® herbicide

In order to provide postemergence control of specified broadleaf weeds, annual and perennial grass weeds and yellow nutsedge, apply a combination of **Biscayne** and Segment. Make applications as a directed spray. Avoid ornamental plant foliage. If ornamental foliage or the foliage of a desirable plant comes into contact with the application solution, immediately wash the solution off.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Biscayne plus Tower® herbicide

In order to control specified emerged broadleaf species (listed in this label) and yellow nutsedge, treat with a tank mix of **Biscayne** plus Tower.

This combination will also provide preemergence control of grass and broadleaf species listed in the Tower label.

Make applications as a directed spray. Avoid ornamental plant foliage. If ornamental foliage or the foliage of a desirable plant comes into contact with the application solution, immediately wash the solution off.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Other Tank Mixes

This product may be tank mixed with other compatible products registered for use in ornamentals.

Make applications as a directed spray. Avoid ornamental plant foliage. If ornamental foliage or the foliage of a desirable plant comes into contact with the application solution, immediately wash the solution off.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When making tank mixes other than those specified in this label, test the tank mix in a small area and allow 5 - 7 days to ascertain potential damage prior to wider application.



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not store at less than 32°F and do not allow product to freeze. Do not store or use near oxidizing agents.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, incineration, or other procedures allowed by state and local authorities.

**Nonrefillable Container (greater than five gallons):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, incineration, or other procedures allowed by state and local authorities.

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

**Biscayne™** is a trademark of Atticus, LLC.

Buctril and Liberty are registered trademarks of Bayer CropScience.

Clarity, Distinct, Facet, Lightning, Outlook, Paramount, Poast, Poast Plus, Pursuit, Raptor, Scepter, Segment and Tower are registered trademarks of BASF.

Classic and Synchrony are registered trademarks of E.I. du Pont de Nemours and Co.

Cobra and Resource are registered trademarks of Valent USA Corp.

FirstRate, Lorsban, Stinger and Turflon are registered trademarks of Dow AgroSciences LLC.

Flexstar and Reflex are registered trademarks of a Syngenta Group Company.

Image is a registered trademark of Central Garden & Pet Company.

Londax, Storm and Ultra Blazer are registered trademarks of United Phosphorus, Inc.

Pounce is a registered trademark of the FMC Corporation.

Roundup is a registered trademark of Monsanto.

20190304a