
SUPPLEMENTAL LABELING

Syngenta Crop Protection LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1162A-S1 0913

GROUP	14	HERBICIDE
--------------	-----------	------------------

Boundary® 6.5EC Herbicide

This supplemental label expires on May 15, 2016 and must not be used or distributed after this date.

Active Ingredients:	
S-metolachlor*:	58.2%
Metribuzin**:	13.8%
<hr/>	
Other Ingredients*:	28.0%
Total:	100.0%

*CAS No. 87392-12-9

**CAS No. 21087-64-9

Contains 5.25 lbs. of S-metolachlor and 1.25 lbs. of metribuzin per gallon.

*Contains approximately 14% petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. 100-1162

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Boundary 6.5EC Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Table 1: Crop Rotation Intervals^{1,3}

Rotational Interval After Application of Boundary 6.5EC²				
4 months	4½ Months	8 Months	12 Months	18 Months
Corn	Winter Barley Winter Wheat Alfalfa	Peas Rice Spring Barley Spring Wheat	Asparagus Cotton Forage Grasses Lentils Sainfoin Sugarcane Tomatoes Other Crops not listed (except root crops)	Onions Sugar Beets and Other Root Crops

¹Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. Stand reductions may occur in some areas.

²Crop rotation recommendations do not include restrictions for the tank mix partner. Refer to the label of the other product for additional restrictions.

³Refer to the specific crop use sections for additional crop rotation precautions.

Postemergence Directed Application (AR, LA, MO – Bootheel only, MS, TN)

Boundary 6.5EC can be applied postemergence directed to soybeans to provide residual control of weeds that emerge after crop emergence in the states of Arkansas, Louisiana, Missouri - Bootheel only, Mississippi and Tennessee. A postemergence directed spray of Boundary 6.5EC can be applied to soybeans in addition to a preemergence or preplant application of Boundary 6.5EC according to label directions. The total amount of Boundary 6.5EC applied must not exceed 3.9 pints per acre per season.

See the table below for Boundary 6.5EC postemergence directed rates according to soil type and organic matter level.

Soil Texture	Broadcast Rate Per Acre	
	0.5% to less than 3% organic matter	3% organic matter or greater
Coarse Loamy sand, sandy loam (over 2% organic matter)	1.3 pt.	1.5 pt.
Medium	1.5-2.0 pt.	2 pt.
Fine	2 pt.	2 pt.
Mississippi Delta only Silty clay, clay	2 pt.	2 pt.

A postemergence directed application of Boundary will provide residual preemergence weed control of the weeds listed in Table 5.

Apply in 10 to 20 gallons of water per acre in a 6 to 8 inch band on each side of the row when soybeans are at least 8 inches tall. Do not allow the directed spray to contact more than the lower $\frac{1}{4}$ to $\frac{1}{3}$ of soybean plants. Soybean leaves contacted by the spray will be killed or severely injured. Do not apply directly to soybeans or serious injury will occur.

Precautions: If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

Post-Directed Application Tank Mixes - Glyphosate Tolerant Soybeans Only

Postemergence directed applications of Boundary 6.5EC can be tank mixed with glyphosate such as Touchdown or Roundup brands in glyphosate-tolerant soybeans only. Refer to the tank-mix partner label for use directions, restrictions and limitations. The most restrictive product labeling applies.

Postemergence Directed Applications – Restrictions

- Do not exceed a total of 3.9 pints per acre per season of Boundary 6.5EC.
- Do not graze or feed treated soybean forage, hay, or straw to livestock.
- Do not apply within 90 days of soybean harvest.
- Boundary 6.5EC cannot be applied to sandy loam or loamy sand soils with less than 2% organic matter.

Boundary® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2013 Syngenta