

GROUP**M5****FUNGICIDE****FIFRA 24(c) Special Local Need Label (SLN)**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
WISCONSIN**For Control of Diseases on Long-Season Potatoes****BRAVO WEATHER STIK®****For Agricultural or Commercial Use Only****EPA Reg. No. 66222-276****SLN No. WI-230004****Active Ingredient:**

Chlorothalonil (tetrachloroisophalonitrile)..... 54.0%

Other Ingredients..... 46.0%**Total:** 100.0%

Contains 6.0 pounds chlorothalonil per gallon

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling and federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on this Supplemental label and the EPA registered label for EPA Reg. No. 66222-276.

It is a violation of Federal law to harm an endangered species. Use of this product may be in counties which have endangered and threatened species present; therefore, applicators should evaluate the areas to be treated and take necessary precautions to avoid harming endangered species or their habitat/environment. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608-224-4547.

Crop	Diseases (Pathogen)	Pint Product/A (lb ai/A)	Application Directions
Potato (long-season varieties)	Late Blight (<i>Phytophthora infestans</i>)	Low rate: $\frac{3}{4}$ (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early Blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	Higher rates: 1 to 1½ (0.75 to 1.125)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> • Vines close between rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo Weather Stik may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Weather Stik label.</p>
	Black dot (<i>Colletotrichum coccodes</i>)		

Specific Use Directions: Do not apply more than 21 ½ pints Bravo Weather Stik (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo Weather Stik is a registered trademark of ADAMA

24(c) Registrant:
Makhteshim Agan of North America, Inc. d/b/a
ADAMA
8601 Six Forks Road, Suite 300
Raleigh, NC 27615

