



FIFRA 24(c) Special Local Need Label (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH DAKOTA

For Control of Diseases on Long-Season Potatoes

Bravo Weather Stik®

For Agricultural or Commercial Use Only

EPA Reg. No. 66222-276

SLN No. ND-190004

Active Ingredient:

Chlorothalonil (tetrachloroisophalonitrile)54.0%

Other Ingredients46.0%

Total:100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

This label is valid until December 31, 2028 or until otherwise amended, withdrawn, cancelled, or suspended

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling and federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on this Supplemental label and the EPA registered label for EPA Reg. No. 66222-276.

Crop	Diseases (Pathogen)	Pint Product/A (lb ai/A)	Application Directions
Potato (long-season varieties)	Late Blight (<i>Phytophthora infestans</i>) Early Blight (<i>Alternaria solani</i>) Botrytis vine rot (<i>Botrytis cinerea</i>)	Low rate: $\frac{3}{4}$ (0.6) -then- Higher rates: 1 to 1½ (0.75 to 1.125)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days). Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events
Crop	Diseases (Pathogen)	Pint Product/A (lb ai/A)	Application Directions
	Black dot (<i>Colletotrichum coccodes</i>)		occur: <ul style="list-style-type: none">• Vines close between rows• Late blight forecasting measures 18 disease severity values (DSV)• The crop reaches 300 P-days Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. Bravo Weather Stik may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Weather Stik label.

Specific Use Directions: Do not apply more than 21 ½ pints Bravo Weather Stik (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo Weather Stik is a registered trademark of an ADAMA Group Company.

24(c) Registrant:
Makhteshim Agan of North America, Inc. d/b/a ADAMA
8601 Six Forks Road
Suite 300
Raleigh, NC 27615



Rev. 10/19/2023