

**FIFRA 24(c) Special Local Need Label (SLN)**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF  
WISCONSIN**For Control of Disease on Long-Season Potatoes****BRAVO ZN<sup>®</sup>****For Agricultural or Commercial Use Only****EPA Reg. No. 66222-278****SLN No. WI-230002****Active Ingredient:**

Chlorothalonil (tetrachloroisophalonitrile)..... 38.5%

**Other Ingredients**..... 61.5%**Total:** 100.0%

Contains 4.17 pounds chlorothalonil per gallon

**KEEP OUT OF REACH OF CHILDREN****WARNING**

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended

**Directions for Use**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling and federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on this Supplemental label and the EPA registered label for EPA Reg. No. 66222-276.

It is a violation of Federal law to harm an endangered species. Use of this product may be in counties which have endangered and threatened species present; therefore, applicators should evaluate the areas to be treated and take necessary precautions to avoid harming endangered species or their habitat/environment. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608-224-4547.

Crop	Diseases (Pathogen)	Lb Product/A (lb ai/A)	Application Directions
<b>Potato</b> (Long-season varieties)	Late Blight ( <i>Phytophthora infestans</i> )	Low rate: 1 1/8 (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early Blight ( <i>Alternaria solani</i> )	-then-	
	Botrytis vine rot ( <i>Botrytis cinerea</i> )	Higher rates: 1 1/2 to 2 1/4 (0.75 to 1.125)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:
	Black dot ( <i>Colletotrichum coccodes</i> )		<ul style="list-style-type: none"> <li>• Vines close between rows</li> <li>• Late blight forecasting measures 18 disease severity values (DSV)</li> <li>• The crop reaches 300 P-days</li> </ul> <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo ZN may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo ZN label.</p>
<b>Specific Use Restrictions:</b> Do not apply more than 30½ pts Bravo ZN (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.			

Bravo ZN is a registered trademark of ADAMA.

24(c) Registrant:  
Makhteshim Agan of North America, Inc. d/b/a  
ADAMA  
8601 Six Forks Road, Suite 300  
Raleigh, NC 27615

