

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee)

Recommendation

EPA Reg. No. 279-3108

**FOR DISTRIBUTION AND USE
ONLY IN CALIFORNIA**

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for BRIGADE WSB Insecticide/Miticide is valid until 11/16/2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Strawberries (0 Day PHI)	Vinegar Flies (adults) (<i>Drosophila</i> spp. Including Spotted-wing <i>Drosophila</i>)	8 - 32 oz/A

DIRECTIONS FOR USE: Thorough coverage is essential to achieve control. Use a minimum spray volume of 5 gallons by air or 50 gallons by ground to obtain full coverage of foliage. Multiple applications may be required.

RESTRICTIONS:

Do not apply more than 80 oz/A (0.5 lb ai/A) per season. Harvesters and other personnel performing tasks with all-day foliage contact in treated fields within five (5) days of application must wear long-sleeved shirt, long pants, and shoes plus socks.

Following treatment of strawberry fields at rates of BRIGADE WSB Insecticide/Miticide greater than 0.1 lb ai/A, harvesters must wear gloves for five (5) days following application.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000