

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

BRIGADIER[®]
INSECTICIDE

**FIFRA 2(ee)
Recommendation
EPA Reg. No. 279-3332**

**NOT FOR USE IN AK, DC, HI, IN,
NH, NM, NY & TX**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE
AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR
APPROVED BY THE EPA.**

This recommendation for BRIGADIER Insecticide is valid until 03/18/2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Soybean	Kudzu bug or Bean Plataspid	6.1 oz/A
DIRECTION FOR USE: Apply 6.1 oz/A BRIGADIER Insecticide by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Thorough coverage is essential to achieve control		
RESTRICTIONS: <ul style="list-style-type: none">Do not apply more than 6.1 fluid ounces of product per acre per application.Do not apply more than 17.92 fluid ounces of product per acre per season.Do not sequentially apply BRIGADIER Insecticide less than 30 days apart.Do not harvest soybeans within 45 days of a BRIGADIER Insecticide application for feeding of dry vines.Do not harvest soybeans within 18 days of a BRIGADIER Insecticide application for feeding of green vines.		



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000