



**2(ee) Recommendation**

EPA Reg. No. 60063-83

**FOR CONTROL OF TAR SPOT ON CORN (FIELD CORN, CORN GROWN FOR SEED, POPCORN)**

For use in the States of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ AND FOLLOW THIS SECTION 2(ee) AND ENTIRE BRIXEN FUNGIDE LABEL BEFORE USING.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Tar Spot ( <i>Phyllachora maydis</i> )	13 - 19 fl. oz.	<p>Apply this product preventively, before disease outbreak, when conditions are favorable to disease development.</p> <p>Application timing:</p> <ul style="list-style-type: none"> <li>• Early application at V4-V8 corn growth stage.</li> <li>• Late application at VT-R3 corn growth stage.</li> </ul> <p>Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 2 gallons of spray suspension per acre by aircraft.</p> <p>Include this product in an integrated pest management program. Alternate applications with a fungicide with a different mode of action.</p> <p>Do not make more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p>

**RESTRICTIONS**

- Do not apply more than 19 fl. oz. this product (0.09 lb. of tetraconazole active ingredient) per acre per year.
- Do not apply a total of more than 2.0 lbs. of azoxystrobin active ingredient per acre per year.
- Do not make more than two applications per year.
- Do not harvest silage within 21 days of an application.
- Pre-Harvest Interval (PHI): 7 days
- Restricted-entry interval (REI): 12 hours for all activities with the exception of 20 days for detasseling corn grown for seed.
- Do not apply after corn growth stage R3 (milk stage).
- Do not apply with adjuvants in sprays made between V8 (8 leaves with collar visible) and VT (last branch of tassel is completely visible).

**NOTICE**

The directions, restrictions and precautions on this and the entire Brixen Fungicide label must be read and followed carefully. This supplemental bulletin must be in the possession of the user at the time of application. Read the entire label including the **WARRANTY AND LIMITATION OF DAMAGES** prior to use.

2019-07-18 Brixen 2(ee) v2