



2(ee) Recommendation

EPA Reg. No. 60063-83

FOR LOWER RATE ON CORN (FIELD CORN, CORN GROWN FOR SEED, POPCORN) AND SOYBEANS

For use in the States of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ AND FOLLOW THIS SECTION 2(ee) AND ENTIRE BRIXEN FUNGIDE LABEL BEFORE USING.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

CORN Field Corn, Corn Grown for Seed, Popcorn		
Diseases Controlled	Rate per Acre	Application Directions
Anthracnose (<i>Colletotrichum graminicola</i>) Common rust (<i>Puccinia sorghi</i>) Eye spot (<i>Aureobasidium zeae</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>) Northern corn leaf blight (<i>Exserohilum turcicum</i>) Northern corn leaf spot (<i>Bipolaris zeicola</i>) Southern corn leaf blight (<i>Bipolaris maydis</i>) Southern rust (<i>Puccinia polysora</i>) Tar Spot (<i>Phyllachora maydis</i>)	10 – 19 fluid ounces	Apply this product preventively when conditions are not conducive to disease development. Application timing: <ul style="list-style-type: none"> • Early application at V4-V8 corn growth stage. • Late application at VT-R3 corn growth stage. The higher rates in the rate range may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions are favorable for disease development. In case of high risk of disease outbreak make a second application of another fungicide. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.
RESTRICTIONS Do not make more than one application per year Do not apply more than 0.09 lb of tetraconazole a.i./acre per year and 2.0 lb/acre of azoxystrobin a.i./acre per year Do not apply after corn growth stage R3 (milk stage) Do not apply with adjuvants in sprays made between V8 (8 leaves with collar visible) and VT (Last branch of tassel is completely visible) Do not apply within 7 days of harvest (7 day PHI). Do not harvest silage within 21 days of an application.		

Soybean		
Diseases Controlled	Rate per Acre	Application Directions
Anthracnose (<i>Colletotrichum</i> spp.) Brown Spot (<i>Septoria glycines</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Powdery Mildew (<i>Microsphaera diffusa</i>) Purple Seed Stain (<i>Cercospora kikuchii</i>) White Mold/Sclerotinia Stem Rot (<i>Sclerotinia sclerotiorum</i>) Asian Soybean Rust (<i>Phakopsora pachyrhizi</i>) Aerial Blight (<i>Rhizoctonia solani</i>)	10 – 16 fluid ounces	<p>Apply this product preventively when conditions are not conducive to disease development.</p> <p>Application timing:</p> <ul style="list-style-type: none"> • Early application at early pod fill growth stage. • Late application at R5 soybean growth stage. <p>The higher rates in the rate range may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions are favorable for disease development.</p> <p>In case of high risk of disease outbreak make a second application at 16 fl oz/acre.</p> <p>Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 2 gallons of spray suspension per acre by aircraft.</p>
RESTRICTIONS Do not apply more than 29.2 fl. oz. of this product per acre per year. Do not make more than one application to soybean forage and hay. Do not make more than 2 applications per acre per year, the total amount applied must not exceed 29.2 fl.oz. of this product (0.137 lbs. a.i./A). Maximum total amount of tetraconazole active ingredient (lbs. a.i./A) which may be applied from all products per acre per year: 0.137 lbs. a.i./A. Maximum total amount of azoxystrobin active ingredient (lbs. a.i./A) which may be applied from all products per acre per year: 1.5 lbs. a.i./A. Do not apply after R5 stage (beginning seed). Do not graze or feed treated forage, silage or hay to livestock. Do not harvest immature soybeans for consumption after plants have been treated. Do not apply to vegetable soybeans grown for immature pods. Pre-Harvest Interval (PHI): 14 days		

NOTICE

The directions, restrictions and precautions on this and the entire Brixen Fungicide label must be read and followed carefully. This supplemental bulletin must be in the possession of the user at the time of application. Read the entire label including the **WARRANTY AND LIMITATION OF DAMAGES** prior to use.

2022-07-20 Brixen 2(ee)

Expiration Date: December 31, 2025