



CGS-10

CGS-10 Controls Russet Mites, Broad Mites, Spider Mites plus Their Eggs, Thrips, Aphids, Nematodes & Other Listed Pests and Diseases. CGS-10 can be applied as a foliar spray or a drench to soil for Burrowing nematodes or soil diseases.

DIRECTIONS FOR USE

Add water to container. **Shake CGS-10 REALLY WELL** before measuring. Dilute **CGS-10** with water at the rate necessary for application. Keep Solutions thoroughly agitated during application. Spray sufficient amount to obtain uniform plant coverage. Certain crops may be sensitive to the final spray solution, test spray a few plants to assess sensitivity. If dilution ratio is modified, also test spray to assess crop sensitivity. Re-treatment may be necessary to achieve and/or maintain control during periods of high pest pressure, or signs of renewed pest activity.

***Important*: Spray early morning or late evening. Never apply during times of peak sun exposure.**

FOR USE ON ALL CROPS - FROM SEEDLINGS TO HARVEST

PRECAUTIONARY STATEMENTS - ENVIRONMENTAL HAZARDS

This product is harmful to bees exposed to direct treatment on blooming - nectar producing crops. Do not apply this product if bees may forage on the plants during this period.

WARNING / AVISO

If you do not understand the label instruction, find someone to explain them to you in detail or call the number provided.

Si usted no entiende las instrucciones impresas en la etiqueta, pídale a alguien que se las explique en forma detallada.

Warranty: Any liability express or implied is limited to purchase of this product.

Caution:

Keep out of reach of children & animals. Harmful if swallowed. Avoid contact with eyes, skin or clothing.

First Aid:

If in eyes, flush with water for at least 15 minutes. If on skin, wash with plenty of soap and water. Get medical attention irritation persists.

Storage and Disposal: Store only in original container in a cool, dry place inaccessible to children and pets. Do not reuse empty container. Offer for recycling if available or discard in trash.

APPLICATIONS

Pest / Diseases Controlled	Maximum Number of Application per week	Minimum Days To Wait before Reapplying	Minimum Days To Wait From Last Application To Harvest
Broad / Russet Mites	3	1	1
Spider Mites	3	1	1
Aphids	3	2	1
Thrips	3	2	1
White Flies	2	2	1
Powdery Mildew	2	2	1
Fungus Gnats	2	2	1
Nematodes	2	3	1
Pythium	2	3	1
Root Rot	2	3	1

**PESTICIDE
INSECTICIDE
MITICIDE
FUNGICIDE**

**PROTECT YOUR CROPS
& INVESTMENT**

Active Ingredients:

Soybean oil* (8001-22-7)..... 39%

Sodium Lauryl Sulfate (151-21-3) 19%

Inert Ingredients: 42%

Water, Isopropyl Alcohol, Sodium Citrate, Citric Acid

*Soybean oil grown organically

This product is exempt from the requirement of a US EPA registration number under FIFRA Section 25(b) as outlined in title 40CFR Section 152.25

HOW TO MIX

Infestation Pressure	Rate of CGS-10 per		
	Gallon (128 fl oz) Spray	100 Gallon Tank	Acre* 43,600 sq. ft.
Normal	1.28 oz	1 gal (128 fl.oz)	.78 gal (100 fl.oz)
Heavy	2 oz	1.56 gal (200 fl. oz)	1.56 gal (200 fl. oz)
Preventative / Maintenance	1/2 oz	0.39 gal (50 fl. oz)	0.39 gal (50 fl. oz)

*Coverage per acre will vary between crops, plant density per acre and canopy size.

FOR BURROWING NEMATODES, SOIL PESTS & PATHOGENS: Drench soil around root zone with normal rate solution until desired coverage is achieved. Application will vary between crop and soil type. For heavy infestations, use heavy solution once a week and drench more frequently.

Manufactured by:

Central Coast Garden Products Inc.
1354 Dayton St Unit N Salinas Ca 93901
Tel: 831-757-4733

www.centralcoastgarden.com

Net Wt: 1 gallon (3.78 Liters)